

GROUP 1 FUNGICIDE

PULL HERE TO OPEN ►



# Mertect<sup>®</sup> 340-F

**syngenta<sup>®</sup>**

## Fungicide

*Active Ingredient:*

Thiabendazole\* ..... 42.3%

*Other Ingredients:* ..... 57.7%

*Total:* ..... 100.0%

\*CAS No.148-79-8

Mertect 340-F is a suspension concentrate containing 4.1 pounds of Thiabendazole per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

See additional precautionary statements and directions for use inside booklet.

**EPA Reg. No. 100-889    EPA Est. 39578-TX-1**

**Product of India**

**Formulated in the USA**

**SCP 889A-L1T 0916  
4073708**

## 2 gallons

Net Contents

®

| FIRST AID   |  |
|---|--|
| <b>If Inhaled:</b>  | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul> |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment.  |  |
| <p align="center"><b>HOT LINE NUMBER</b><br/> For 24-Hour Medical Emergency Assistance (Human or Animal) or<br/> Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),<br/> Call<br/> <b>1-800-888-8372</b></p> |  |

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment (PPE)

**Mixers, loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

In addition to the above, handlers making application to mushroom houses using hand held sprayers must wear chemical-resistant gloves.

### User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

*continued...*

## **PRECAUTIONARY STATEMENTS (*continued*)**

### **Environmental Hazards**

This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading.

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves: barrier laminate, butyl rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton®  $\geq 14$  mils
- Shoes plus socks

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.**

## PRODUCT INFORMATION

Mertect 340-F is a systemic fungicide which aids in the control of many important plant diseases. If treatment is not effective following use of Mertect 340-F as recommended, a tolerant strain of fungus may be present. Consideration should be given to prompt use of other suitable non-benzimidazole fungicides. Mertect 340-F is not for use in formulating agricultural products.

## RESISTANCE MANAGEMENT

|       |   |           |
|-------|---|-----------|
| GROUP | 1 | FUNGICIDE |
|-------|---|-----------|

This product contains Thiabendazole, a Group 1 fungicide. Thiabendazole belongs to the methylbenzimidazole carbamate class of chemistry which disrupts  $\beta$ -tubuline assembly in mitosis.

Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include alternation and/or tank-mixing with products having different modes of action or limiting the total number of applications per season.

Syngenta encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

### For additional information on Fungicide Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481
- Contact your local extension specialist or certified crop advisor
- Visit the Fungicide Resistance Action Committee (FRAC) on the web at:  
<http://www.frac.info>

## CROP USE DIRECTIONS

GENERAL - SHAKE OR STIR WELL BEFORE USE.

Clean equipment before using Mertect 340-F. **When Mertect 340-F is dispersed in water, the resulting suspension must be constantly agitated.** Avoid foaming.

Use the higher specified rates under conditions of heavier disease pressure.

Compatibility information for Mertect 340-F with other chemicals is limited. Consult your supplier or agricultural experiment station for specific recommendations.

**Carrots - *Botrytis cinerea* Gray Mold and *Sclerotinia sclerotiorum* (Sclerotinia Rot):** Dip carrots, before storage, for 5-10 seconds in a suspension containing 41 fl oz of Mertect 340-F per 100 gal of water. Replenish the suspension when the volume is too low or when it becomes dirty.

The runoff and wastes from the dipping operation should not be discarded in a drainage which could contaminate public water systems.

**Mushrooms (*Agaricus* spp.) - Dry Bubble (*Verticillium fungicola*), Wet Bubble (*Mycogone perniciosa*), Cobweb (*Dactylium dendroides*) and Green Mold (*Trichoderma* spp.):**

FOR USE AS AN ADMIX WITH MUSHROOM SPAWN GRAINS, MUSHROOM GROWING SUPPLEMENT, AND/OR AS A DRENCH DURING NORMAL WATERING OPERATIONS FOR MUSHROOM GROWING BEDS AT CASING OR BETWEEN BREAKS

**NOTE:** Do not exceed a combined total of 20 fl oz per 1,000 sq ft of product, between the various applications to spawn grains, supplement, and mushrooms. This amount may be used in various treatment regimes. Please refer to the tables below for the rates for the various usage so that the 20 fl oz per 1,000 sq ft is not exceeded.

**Spawn Grains:** Apply a maximum of 5.4 fl oz of Mertect 340-F per 1,000 sq ft. Follow the table below to determine the fl oz of Mertect 340-F that can be mixed with the spawn to deliver 5.4 fl oz per 1,000 sq ft.

#### **Dosage Table for Use on Spawn Grains**

Do not apply Mertect 340-F to gypsum, limestone, or chalk. Admix Mertect 340-F directly onto spawn grain.

| Pounds spawn per 8000 sq ft | Max. fl oz Mertect per 100 pounds spawn |
|-----------------------------|---|
| 960                         | 4.5                                     |
| 1040                        | 4.1                                     |
| 1120                        | 3.8                                     |
| 1200                        | 3.6                                     |
| 1280                        | 3.4                                     |
| 1360                        | 3.2                                     |
| 1440                        | 3.0                                     |

**Mushroom Supplements:** A finished batch of mushroom supplement can contain up to 0.1% Thiabendazole (by weight). The Thiabendazole added to the supplement must be included as part of the total of the 20 fl oz of Mertect 340-F allowed per 1,000 sq ft per crop. The total amount of Mertect 340-F that can be applied from all treatments per crop is dependent on the amount of supplement applied (use the table below to determine the remaining volume of Mertect 340-F that can be applied per crop).

| Pounds supplement* per 8,000 sq ft | Fl oz Mertect 340-F remaining that may be applied to mushrooms, supplement, casing, or other timing | Fl oz Mertect 340-F remaining that may be applied to mushrooms, supplement, casing, or other timing if a spawn treatment was also applied |
|------------------------------------|---|---|
| 800                                | 17  | 11.6  |
| 1200                               | 15  | 9.6   |
| 1400                               | 14.5  | 9.1   |
| 1800                               | 13  | 7.6   |
| 2200                               | 11  | 5.6   |
| 2400                               | 9   | 3.6   |

\*Assumes at maximum rate of 0.1% Thiabendazole.

The amount added for supplement treatment must be deducted from the total remaining amount that can be applied to mushrooms so as not to exceed 20 fl oz per 1,000 sq ft per crop.

**Note:** If you are unsure if the mushroom supplement that you are using contains Thiabendazole, contact the manufacturer of the mushroom supplement.

**Applying to Mushrooms:** Refer to the tables above for the amount of Mertect 340-F used in any spawn grain or supplement treatments before applying Mertect 340-F to the mushrooms to ensure that 20 fl oz per 1,000 sq ft is not exceeded.

Casing Applications: Apply no more than 8 fl oz Mertect 340-F per 1000 sq ft. Mertect 340-F may be applied at casing as a drench during normal watering operations.

Other Timings: Apply no more than 4 fl oz Mertect 340-F per 1000 sq ft.

Do not apply within 12 hours prior to harvest.

**Ornamental Bulbs and Corms (One Application Only) - *Fusarium* Basal Rot and *Penicillium* Blue Mold:** Clean and treat bulbs or corm within 24-48 hours of digging. Prepare suspension of 30 fl oz of Mertect 340-F per 100 gal water. Submerge stock completely in dipping suspension. Suspension temperature should be 55-75°F (13-24°C). Discard suspension (1) when it becomes dirty, (2) after using five times, or (3) after 24 hours, whichever occurs first.

**Fusarium (Bulbs and Corms):** Dip bulbs 15-30 minutes, corms 15 minutes.

**Blue Mold (Bulbs):** Dip 10-15 minutes.

After treatment, dry bulbs or corms in a shaded, well-ventilated area. Curing or retarding may precede or follow treatment.

**Potato - *Fusarium* Tuber Rot:** Mist unwashed tubers on a conveyor line, with tumbling action, entering storage with 0.42 fl oz of Mertect 340-F to each 2,000 lb of tubers in sufficient water for complete coverage. If an additional treatment is necessary before shipping, mist the tubers at the same rate or dip the tubers for 20 seconds in a solution containing 0.42 fl oz of Mertect 340-F per gal of water. Do not treat seed potatoes after cutting.

Mertect 340-F for potatoes should be used as part of an Integrated Management Program, which includes prevention of bruising during harvest and maintaining proper storage temperature and relative humidity. Neither cultural practices nor chemical measures will prevent *Fusarium* infection if potatoes are damaged or improperly stored.

**Pome Fruit (Apples and Pears) - Blue Mold Rot, Bull's Eye Rot, and Gray Mold (Nest Rot, Cluster Rot), Stem End and Neck Rot:** Dip, flood, or spray harvested fruit with a suspension of 16 fl oz of Mertect 340-F in 100 gal of water. Do not treat for over 3 minutes. Treat apples only before and after storage for maximum decay control. Treat pears only once.

**Sweet Potato (One Application Only) "Seed Roots" - Black Rot, Scurf, and Foot Rot:** Dip the seed roots in a suspension containing 8 fl oz of Mertect 340-F per 7.5 gal of water (107 fl oz per 100 gal). Treat the seed roots for 1-2 minutes and plant immediately. Discard the suspension when the volume is too low or when it becomes dirty. Do not use the treated roots for food or feed.

## SEED TREATMENT

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

Seed treatment with Mertect 340-F provides protection from seed and soil-borne pests listed under target diseases, thereby reducing the likelihood of encountering early-season plant infection, or late-season disease symptoms resulting from early-season infection.

For protection against other seed and soil-borne diseases not controlled by Mertect 340-F, it is highly recommended that a compatible seed treatment fungicide be used at EPA registered label rates and tank mixed in combination with Mertect 340-F. These fungicides must show safety on treated seed, alone or in combination with Mertect 340-F. Consult the label for appropriate use rates and follow all use instructions.

## **MIXING PROCEDURES**

Apply Mertect 340-F as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the recommended amount of Mertect 340-F into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment.

- Use an EPA-approved dye or colorant that imparts an unnatural color to the seed as required in 40 CFR 153.155 (c).
- Allow seed to dry before bagging.

### **Tank Mixtures**

Confirm physical compatibility, seed safety, seed safety in stored seed, and biological efficacy before tank mixing with other seed treatment products.

Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibiting such mixing.

## **SEED CONTAINER LABEL REQUIREMENTS**

Federal law requires that bags containing treated seed shall be labeled with the following information:

- This seed has been treated with Thiabendazole fungicide.
- Do not use for feed, food, or oil purposes.

In addition, include the following statements on bags containing seeds treated with Mertect 340-F:

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.



- In the event of crop failure or after harvest of crops grown from seed treated with Mertect 340-F, the following crops may be replanted immediately:

| Immediate Plantback   |
|---|
| Alfalfa   |
| Cereal Grains: Barley, Corn, Oat, Rye, Triticale, and Wheat |
| Cucurbit Vegetables<br>Crop Group 9                         |
| Head and Stem <i>Brassica</i><br>Crop Subgroup 5A           |
| Legume Vegetables (Succulent or Dried)<br>Crop Group 6      |
| Onion, Bulb,<br>Crop Subgroup 3-07A                         |
| Root Vegetables (Except Sugarbeet)<br>Crop Subgroup 1B      |
| Spinach   |
| Sweet Potato  |

- For any other crop, the minimum plant back interval is 30 days from the date seed treated with Mertect 340-F was planted.
- Use crop-specific seeding rates according to local agricultural practice.
- Regardless of type of application, do not apply more than 0.15 lb Thiabendazole per acre (68.0 grams ai/A) per year

Use an EPA-approved dye or colorant that imparts an unnatural color to the seed as required in 40 CFR 153.155 (c).

| Crop                           | Target Diseases   | Use Rate<br>mg ai/seed | Use Rate<br>fl oz product/<br>100 lb seed | Remarks   |
|--------------------------------|---|------------------------|---|---|
| Alfalfa                        | Seedling diseases caused by <i>Fusarium</i> spp.              | 0.004                  | 5.8                                       | Fl oz product/100 lb seed calculations assume an average seed size of 210,000 seeds/lb.   |
| Cereals, small grain<br>Barley | Seedling diseases caused by <i>Fusarium</i> spp.              | 0.03                   | 2.6                                       | Fl oz product/100 lb seed calculations assume an average seed size of 12,500 seeds/lb.  |
| Oats                           | Seedling diseases caused by <i>Fusarium</i> spp.              | 0.03                   | 3.4                                       | Fl oz product/100 lb seed calculations assume an average seed size of 16,400 seeds/lb.  |
| Rye                            | Seedling diseases caused by <i>Fusarium</i> spp.              | 0.03                   | 3.2                                       | Fl oz product/100 lb seed calculations assume an average seed size of 15,600 seeds/lb.  |
| Triticale                      | Seedling diseases caused by <i>Fusarium</i> spp.              | 0.03                   | 2.6                                       | Fl oz product/100 lb seed calculations assume an average seed size of 12,750 seeds/lb.  |
| Wheat, Winter and Spring       | Seed-Borne Fusarium Scab only                                 | 0.0016                 | 0.17                                      | Fl oz product/100 lb seed calculations assume an average seed size of 15,000 seeds/lb*. For best control of seed-borne Fusarium scab Mertect 340-F should be combined with Maxim 4FS, a pre-mix product containing fludioxonil, or another product with seed-borne Fusarium activity. |
|                                | Seed-Borne Common Bunt (Stinking Smut)                        | 0.013                  | 1.3                                       |   |
|                                | Soil-Borne Common Bunt (Stinking Smut)                        | 0.025                  | 2.6                                       |   |
|                                | Seedling diseases caused by <i>Fusarium</i> spp. (soil-borne) | 0.0195 - 0.039         | 1.95 - 3.9                                |   |
| Winter Wheat Only              | Dwarf Bunt  | 0.025 - 0.039          | 2.6 – 3.9                                 | For normal or late winter wheat plantings. Use the lower rate on late winter wheat plantings only. For best control of Dwarf Bunt Mertect 340-F should be combined with a pre-mix product containing difenoconazole.  |

\*For protection against other seed and soil-borne diseases, it is highly recommended that a compatible seed treatment fungicide such as Dividend Extreme®, Vibrance® Extreme, or CruiserMaxx® Vibrance® Cereals be used at EPA registered label rates and tank mixed in combination with Mertect 340-F. These fungicides must show safety on treated seed, alone or in combination with Mertect 340-F. Consult the label for appropriate use rates and follow all use instructions.

| Crop   | Target Diseases   | mg ai/seed |
|--|---|------------|
| <b>Brassica</b><br>Head and Stem subgroup<br>Broccoli<br>Broccoli, Chinese<br>Brussels sprouts<br>Cabbage<br>Cabbage, Chinese (napa)<br>Cabbage, Chinese mustard<br>Cauliflower<br>Cavalo broccolo<br>Kohlrabi                                       | Seedling diseases caused by <i>Fusarium</i> spp.  | 0.002      |
| <b>Bulb Vegetables</b><br>Daylily<br>Fritillaria<br>Garlic, great-headed<br>Garlic, serpent<br>Lily<br>Onion, Chinese<br>Onion, dry bulb<br>Onion, pearl<br>Onion, potato<br>Shallot<br><br>(Including cultivars, varieties and/or hybrids of these) | Seedling diseases caused by <i>Fusarium</i> spp.  | 0.002      |
| <b>Cucurbit Vegetables</b><br>Chayote (fruit)<br>Chinese waxgourd<br>Citron melon<br>Cucumber<br>Gherkin<br>Gourd<br>Momoridica spp.<br>Muskmelon<br>Pumpkin<br>Summer squash<br>Winter squash<br>Watermelon   | Seedling diseases caused by <i>Fusarium</i> spp.  | 0.1        |
| <b>Spinach</b><br>Baby<br>Mature   | Seedling diseases caused by <i>Fusarium</i> spp.<br><br>Verticillium wilt caused by seed borne <i>Verticillium dahlia</i> | 0.006      |

| Crop  | Target Diseases                                  | mg ai/seed |
|---|--|------------|
| <b>Vegetables, root subgroup (except sugarbeets)</b><br>Beet, garden<br>Burdock, edible<br>Carrots<br>Celeriac<br>Chervil, turnip-rooted<br>Chicory<br>Ginseng<br>Horseradish<br>Parsley, turnip-rooted<br>Parsnip<br>Radish<br>Radish, oriental<br>Rutabaga<br>Salsify, Spanish<br>Skirret<br>Turnip | Seedling diseases caused by <i>Fusarium</i> spp. | 0.002      |

| Crop  | Target Diseases  | mg ai/seed    | Use Rate fl oz product/<br>100 lb seed<br>(g ai/100 kg) |
|---|--|---------------|---|
| <b>Legume Vegetables (Succulent or Dried) Crop Group 6<sup>1</sup></b><br><br><b>BEANS</b><br><br>Bean ( <i>Lupinus</i> spp.): includes grain lupin, sweet lupin, white lupin and white sweet lupin<br><br>Bean ( <i>Phaseolus</i> spp.): includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean and wax bean<br><br>Bean ( <i>Vigna</i> spp.): includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean and yardlong bean<br><br>Broad bean (fava bean) ( <i>Vicia faba</i> )<br><br>Guar ( <i>Cyamopsis tetragonoloba</i> )<br><br>Jackbean ( <i>Canavalia ensiformis</i> )<br><br>Lablab bean (hyacinth bean) ( <i>Lablab purpureus</i> )<br><br>Sword bean ( <i>Canavalia gladiata</i> ) | Seed decay, seedling wilt, and damping-off caused by <i>Phomopsis</i> spp.<br><br>Seedling diseases caused by <i>Fusarium</i> spp. | 0.035 – 0.079 | 0.30 – 0.68 <sup>4</sup><br>(10 – 22)                   |

continued...

| Crop   | Target Diseases  | mg ai/seed | Use Rate fl oz product/<br>100 lb seed<br>(g ai/100 kg) |
|--|--|------------|---|
| <b>CHICKPEA</b><br>Chickpea (Garbanzo Bean)<br>( <i>Cicer arietinum</i> )  | Seed-borne <i>Ascochyta</i> and <i>Phoma</i> spp.<br>which cause Ascochyta Blight (foot rot,<br>basal stem rot or black stem) <sup>2</sup><br><br>Seedling diseases caused by <i>Fusarium</i> spp. | 0.237      | 2.04 <sup>4</sup><br>(65)                               |
| <b>LENTIL</b><br>Lentil ( <i>Lens esculenta</i> )  | Seed-borne <i>Ascochyta</i> and <i>Phoma</i> spp.<br>which cause Ascochyta Blight (foot rot,<br>basal stem rot or black stem) <sup>2</sup><br><br>Seedling diseases caused by <i>Fusarium</i> spp. | 0.014      | 1.05 <sup>4</sup><br>(34)                               |
| <b>PEAS<sup>3</sup></b><br>Pea ( <i>Pisum</i> spp.): includes<br>dwarf pea, edible-pod pea,<br>English pea, field pea, garden<br>pea, green pea, snow pea and<br>sugar snap pea<br><br>Pigeon Pea ( <i>Cajanus cajan</i> )   | Seed-borne <i>Ascochyta</i> and <i>Phoma</i> spp.<br>which cause Ascochyta Blight (foot rot,<br>basal stem rot or black stem) <sup>2</sup><br><br>Seedling diseases caused by <i>Fusarium</i> spp. | 0.077      | 1.02 <sup>4</sup><br>(33)                               |
| <p>1. Untreated foliage of legume vegetables from seeds treated with this product are covered by an EPA tolerance.</p> <p>2. Will not control disease in aerial portions of the plant after emergence due to infection from such as soil or wind. For best results against Ascochyta blight, plant legume vegetable seeds treated with Mertect 340-F fungicide as late in the spring as possible.</p> <p>3. For Export Use on Peas: Mertect 340-F may be mixed with other registered seed treatment products in accordance with the importing nation's phytosanitary requirements. Treated seeds must be labeled: "For Export Only – Not to be Sold or Offered for Sale in the United States."</p> <p>4. Conversion from mg ai/seed are based on 1,250 bean seeds per pound, 1,250 chickpea seeds per pound, 10,600 lentil seeds per pound, and 1,930 pea seeds per pound. If the size of the seed to be treated differs greatly from these numbers use the mg ai/seed rate to calculate the fl. oz./100 lb of seed to be applied.</p> |  |            |   |

| Crop    | Target Diseases   | Use Rate<br>g ai/100 kg<br>seed<br>(mg ai/seed) | Use Rate<br>fl oz product/<br>100 lb seed | Remarks   |
|---------|---|---|---|---|
| Soybean | Pod and stem blight<br>( <i>Phomopsis</i> spp.)   | 2.5-5 g/100 kg<br>(0.004-0.008 mg<br>ai/seed)   | 0.08 - 0.16                               | Fl oz product/100 lb seed<br>calculations assume an<br>average seed size of 3,000<br>soybean seeds/lb.<br><br>For protection against other<br>seed and soil-borne diseases,<br>it is recommended that a co-<br>fungicide such as Syngenta<br>Apron® XL + Maxim® 4FS,<br>or an ApronMAXX® brand<br>fungicide, be used at EPA<br>registered label rates and<br>tank mixed in combination<br>with Mertect 340-F. |
|         | <i>Fusarium</i> spp.<br>(ex. <i>Fusarium virguliforme</i> ,<br><i>Fusarium graminearum</i> ,<br><i>Fusarium oxysporum</i> ,<br><i>Fusarium solani</i> ) | 10-20 g/100 kg<br>(0.015– 0.030 mg<br>ai/seed)  | 0.32 - 0.64                               |   |

#### Restrictions and Limitations

- This product must not be tank mixed with any pesticide unless physical compatibility and phytotoxicity tests have been completed and proved to be satisfactory.
- Regardless of type of application, do not apply more than 0.15 lb Thiabendazole per acre (68.0 grams ai/A) per year.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Store Mertect 340-F in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C). Repeated product freezing and thawing should be avoided. Shake or stir product before use.

#### Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

*continued...*

## **STORAGE AND DISPOSAL (*continued*)**

### **Container Handling [less than or equal to 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling [greater than 5 gallons – mini-bulk]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling [greater than 5 gallons - bulk]**

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

Apron®, ApronMAXX®, Maxim®, Mertect®, the ALLIANCE FRAME   
the SYNGENTA Logo and the PURPOSE ICON   
are Trademarks of a Syngenta Group Company

©2016 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 889A-L1T 0916**  
**4073708**



GROUP 1 FUNGICIDE



## Fungicide

Active Ingredient:  
Thiabendazole\*: . . . . . 42.3%  
Other Ingredients: . . . . . 57.7%  
Total: . . . . . 100.0%

\*CAS No. 148-79-8

Mertect 340-F is a suspension concentrate containing 4.1 pounds of Thiabendazole per gallon.

See additional precautionary statements and directions for use inside booklet.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-889

EPA Est. 39578-TX-1

Mertect® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company

©2016 Syngenta

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

SCP 889A-L1T 0916  
4073708

## KEEP OUT OF REACH OF CHILDREN. CAUTION

### FIRST AID

- |             |  |
|-------------|--|
| If Inhaled: | <ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul> |
|-------------|--|

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

### HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or  
Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),  
Call  
1-800-888-8372

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Remove and wash contaminated clothing before reuse.

#### Environmental Hazards

This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Store Mertect 340-F in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C). Repeated product freezing and thawing should be avoided. Shake or stir product before use.

#### Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

#### Container Handling

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

2 gallons  
Net Contents

syngenta®

