

Disease Risk Fungicide Schedules



PLANTING DATE _____	30 DAYS	45 DAYS	60 DAYS	75 DAYS	90 DAYS	105 DAYS	120 DAYS
TRADITIONAL PROGRAM	LEAF SPOT	LEAF SPOT	LEAF SPOT WHITE MOLD LIMB ROT	LEAF SPOT	LEAF SPOT WHITE MOLD LIMB ROT	LEAF SPOT	LEAF SPOT
LOW RISK*	Bravo® 1.0 pt + Alto® 5.5 oz	21 days →	Elatus® 9.5 oz	21 days →	Elatus® 9.5 oz	21 days →	Bravo® 1.0 pt + Alto® 5.5 oz
MODERATE <i>5 spray program</i>	Bravo® 1.0 pt + Alto® 5.5 oz	21 days →	Elatus® 9.5 oz	Bravo® 1.5 pt	Elatus® 9.5 oz	21 days →	Bravo® 1.0 pt + Alto® 5.5 oz
MODERATE <i>6 spray program</i>	Elatus® 7.3 oz	Bravo® 1.0 pt + Alto® 5.5 oz	Elatus® 7.3 oz	Bravo® 1.5 pt	Elatus® 7.3 oz	Bravo® 1.0 pt + Alto® 5.5 oz	Bravo® 1.5 pt
HIGH RISK	Bravo® 1.0 pt + Alto® 5.5 oz	Bravo® 1.5 pt	Elatus® 9.5 oz	Bravo® 1.5 pt	Elatus® 9.5 oz	Bravo® 1.0 pt + Alto® 5.5 oz	Bravo® 1.5 pt
YOUR PROGRAM							

Programs developed through the cooperation of



* In stewardship of FRAC guidelines, Syngenta recommends tank mixing Bravo Weather Stik® fungicide (containing the active ingredient chlorothalonil) with Abound® fungicide (containing the active ingredient azoxystrobin). When planting late-season varieties that have maturities greater than 140 days, such as Georgia-02C, C-99R and York, spray intervals could be stretched to 24 to 25 days depending on rotation and rainfall patterns. Under conditions of higher than normal rainfall or tropical storm conditions, fungicide spray intervals should be reduced and rates increased to coincide with the next most conservative index recommendation.

Under the PeanutRx™ incentive offered by Syngenta, Syngenta brand fungicides are the only fungicides that may be used in your spray program to qualify for Syngenta standard product performance protection.

©2016 Syngenta. Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status. Abound®, Alto®, Bravo®, Bravo Weather Stik®, Elatus®, the Alliance Frame, the Purpose Icon and the Syngenta logo are trademarks of a Syngenta Group Company. Peanut Rx™ is a trademark of the University of Georgia. Thimet® is a trademark of AMVAC Chemical Corporation. Classic® is a trademark of E. I. du Pont de Nemours and Company. FloRun™ and TUFRunner™ are trademarks of Florida Foundation Seed Producers, Inc. All other trademarks or service marks are the property of their respective owners. Use of third party trademarks are for informational use only and are not intended to imply endorsement by or affiliation with trademark owners. Syngenta Customer Center: 1-866-SYNGENTA (796-4368).

GS 406.61000

SLC 7658A 03-2016



Assess Disease Risk in Your Field and Develop a Peanut Rx

This worksheet will lead you through the four-step process of determining your disease risk level in order to customize a Peanut Rx™ for your individual field using the reverse side of this worksheet and with the assistance of your Syngenta representative.

For each of the risk index factors, identify which option best describes the situation for your field and add the index value associated with each choice to obtain your overall disease risk value. This worksheet does not contain all of the varieties included in the 2016 Peanut Rx or the notes that accompany each factor. To view the complete 2016 Peanut Rx, visit the University of Georgia peanut website at www.ugapeanuts.com.



Step 1: Assess Your Disease Risk

Variety Selection				
Variety ¹	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
Bailey ³	10	15		10
Florida-07 ²	10	20		15
Florida Fancy ²	25	20		20
FloRun™ '107 ²	20	25		20
Georgia-06G	10	20		20
Georgia-07W	10	20		15
Georgia-09B ³	20	25		25
Georgia-12Y	5	15		10
Georgia-13M1,2	10	30		25
Georgia-14N1,2,4	10	15		15
Georgia Green	30	20		25
Georgia Greener ³	10	20		20
Tifguard ⁴	10	15		15
TUFRunner™ '297' ^{1,2}	15	25		20
TUFRunner™ '727' ²	20	15		15
TUFRunner™ '511' ^{1,2}	20	30		15

Planting Date				
Peanuts are planted:	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
Prior to May 1	30	0	10	0
May 1 to May 10	15	0	5	0
May 11 to May 31	5	5	0	0
June 1 to June 10	10	10	0	5
After June 10	15	10	0	5

Plant Population (final stand, not seeding rate)				
Plant stand:	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
Less than 3 plants/ft	25	NA	0	NA
3 to 4 plants/ft (3)	10 (15)	NA	0 (0)	NA
More than 4 plants/ft	5	NA	5	NA

At-plant Insecticide				
Insecticide used	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
None	15	NA	NA	NA
Other than Thimet® 20G	15	NA	NA	NA
Thimet 20G	5	NA	NA	NA

Row Pattern				
Peanuts are planted in:	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
Single rows	10	0	5	0
Twin rows	5	0	0	0

Tillage				
Tillage type	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
Conventional	15	10	0	0
Reduced	5	0	5	5

Classic® Herbicide				
Classic usage	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
Classic applied	5	NA	NA	NA
No Classic applied	0	NA	NA	NA

Crop Rotation (with a non-legume crop)				
Years between peanut crop	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
0	NA	25	25	20
1	NA	15	20	15
2	NA	10	10	10
3 or more	NA	5	5	5

Field History				
Have you had a problem controlling these diseases?	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
No	NA	0	0	0
Yes	NA	10	15	10

Irrigation				
Does the field receive irrigation?	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points	
			White Mold	Limb Rot
No	NA	0	0	0
Yes	NA	10	5	10

¹ Adequate research data is not available for all varieties with regards to all diseases. Additional varieties will be included as data to support the assignment of an index value are available.

² High oleic variety

³ Varieties Georgia Greener and Bailey have increased resistance to *Cylindrocladium* black rot (CBR) than do other varieties commonly planted in Georgia.

⁴ Tifguard has excellent resistance to the peanut root-knot nematode.

Step 2: Calculate Your Severity Points

Fill in the following table to calculate your severity points for each of the four major peanut diseases given the 10 determining factors. Total each column to establish your disease index values.

	Spotted Wilt	Leaf Spot	White Mold	Rhizoctonia Limb Rot
Variety				
Planting Date				
Plant Population				
At-plant Insecticide				
Row Pattern				
Tillage				
Classic Herbicide				
Crop Rotation				
Field History				
Irrigation				
Your Total Index Value				

Step 3: Interpret Your Index Values

Once you've calculated your index values, utilize the following information to interpret your risk level situation.

	Spotted Wilt	Leaf Spot	White Mold	Rhizoctonia Limb Rot
Low Risk	≤ 65	10-35	10-25	TBD
Moderate Risk	70-110	40-60	30-50	TBD
High Risk	≥ 115	65-100	55-80	TBD

When tomato spotted wilt virus incidence is high statewide or in your region, even fields with a low risk level may experience significant losses. Consider the following recommendations to reduce your spotted wilt risk level:

- Use less susceptible varieties
- Adjust your planting date
- Consult the complete Peanut Rx for additional options that may also provide limited benefit

Step 4: Develop Your Peanut Rx

Once you have calculated your total risk for each fungal disease, utilize the most conservative fungicide program as your guide for customizing a per-field prescription spray program with the assistance of your Syngenta representative. Syngenta-recommended fungicide spray programs for each risk level are included on the reverse side of this worksheet.

Programs developed through the cooperation of

