

DIFENOCONAZOLE GROUP 3 FUNGICIDE

FLUDIOXONIL GROUP 12 FUNGICIDE

PULL HERE TO OPEN ►



# Academy™

syngenta®

## Fungicide

*Active Ingredient:*

Difenoconazole:\* ..... 20.9%

Fludioxonil:\*\* ..... 12.5%

*Other Ingredients:* ..... 66.6%

*Total:* ..... 100.0%

\*CAS No. 119446-68-3

\*\*CAS No. 131341-86-1

Academy is a flowable suspension concentrate.

Academy contains 2.06 pounds of difenoconazole and 1.23 pounds of fludioxonil per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1529

EPA Est. 100-NE-001

SCP 1529A-L1A 0222

4170570

## 2.5 gallons

Net Contents

®

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have a person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>HOTLINE NUMBER</b> For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call <b>1-800-888-8372</b>	

<b>PRECAUTIONARY STATEMENTS</b>	
<b>Hazards to Humans and Domestic Animals</b>	<b>CAUTION</b>
Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.	
<b>Personal Protective Equipment (PPE)</b>	
<b>Applicators and other handlers must wear:</b>	
<ul style="list-style-type: none"> <li>• Long-sleeved shirt and long pants</li> <li>• Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, Viton™ ≥ 14 mils</li> <li>• Shoes plus socks</li> <li>• Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N*, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N*, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters</li> </ul>	
<b>Environmental Hazards</b>	
This product is toxic to fish, mammals, aquatic invertebrates, oysters, and shrimp. Do not contaminate water when disposing of equipment washwater or rinsate. DO NOT allow contaminated waste water from the processing areas to enter lakes, streams, ponds or other waters. Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.	
<b>Physical or Chemical Hazards</b>	
Do not use or store near heat or open flame. Do not use or mix with oxidizing agents.	

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

**Do not formulate this product into other end-use products.**

## PRODUCT INFORMATION

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL.**

Academy is a protective fungicide used to aid in the control of several post-harvest diseases **in post-harvest treatment facilities for pome fruit.**

## RESISTANCE MANAGEMENT

DIFENOCONAZOLE	GROUP	3	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

Academy contains two fungicides with different modes of action. Difenoconazole is a broad spectrum fungicide in Group 3. Fludioxonil is a phenylpyrrole in Group 12. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or state agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Syngenta Crop Protection encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label. Academy should not be alternated or tank-mixed with any fungicide in FRAC Group 3 or 12 to which resistance has already developed.

Use sanitation and other cultural practices to minimize disease in order to control disease and prevent or delay disease development.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of difenoconazole or other Group 3 fungicides and fludioxonil or other Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crop and pathogens.
- For further information or to report suspected resistance, contact Syngenta at 1-866-Syngent(a) (866-796-4368). You can also contact your pesticide distributor or university extension specialist to report resistance.

## MIXING PROCEDURES

Vigorously shake the product container before mixing. Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use.

To determine the physical compatibility of Academy with other products, use a jar test as described below.

**Jar Compatibility Test:** Using a quart jar, add the proportionate amounts of the products to 1 qt. of water or wax/oil emulsion. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

If using Academy in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate may be exceeded and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product which prohibits such mixing. Tank mixtures are permitted only in those states where the tank mix partner is registered.

THE CROP SAFETY OF ALL POTENTIAL TANK MIXES INCLUDING ADDITIVES AND OTHER PESTICIDES ON ALL CROPS HAS NOT BEEN TESTED. BEFORE APPLYING ANY TANK MIXTURE, THE SAFETY TO THE TARGET CROP SHOULD BE CONFIRMED.

Add 1/2 of the required amount of water or wax/oil emulsion (or aqueous dilution of a wax/oil emulsion) to the spray or mixing tank. With the agitator running, open the container and add the Academy to the tank. Continue agitation while adding the remainder of the carrier. Begin application of the solution after the Academy has completely and uniformly dispersed into the mix carrier. Maintain agitation until all of the mixture has been applied.

If tank-mixing, add the desired amount of other products recommended for tank mixture after Academy has completely and uniformly dispersed into the mix carrier. Add tank mix partners in this order unless label directions or other considerations indicate otherwise: wettable powders, wettable granules (dry flowables), liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied. Maintain agitation until all of the mixture has been applied.

## CROP USE DIRECTIONS

### Pome Fruit

#### Crop Group 11-10

Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear; Pear, Asian; Quince; Quince, Chinese; Quince, Japanese; Tejocote; and cultivars, varieties and/or hybrids of these.

Use Academy as a post-harvest dip, drench, flood, or spray for the control of post-harvest diseases caused by:

- Alternaria rot (side rot) and surface mold (*Alternaria alternata*)
- Bitter rot (*Colletotrichum gloeosporioides*)
- Blue mold (*Penicillium expansum*)
- Bull's-eye rot (*Neofabraea malacorticis*; *N. alba*; *N. perrenans*; *N. nova*)
- Gray mold (*Botrytis cinerea*)
- Phacidiopycnis rot (*Phacidiopycnis piri*)
- Rhizopus rot (*Rhizopus stolonifer*)
- Speck rot (*Phacidiopycnis washingtonensis*)
- Sphaeropsis rot (*Sphaeropsis pyriputrescens*)
- White rot (*Botryosphaeria dothidea*)

Application Method	Disease	Rate (fl oz)	Remarks
Bin/Truck Drench or In-Line Dip/Drench or Flooder	Alternaria rot and surface mold Bitter rot Blue mold Bull's-eye rot Gray mold Phacidiopycnis rot Rhizopus rot Speck rot Sphaeropsis rot White rot	16 fl oz/100 gallons of water	<ul style="list-style-type: none"> <li>• Ensure proper coverage of the fruit.</li> <li>• For re-cycling in-line drench or dip treatments, the fungicide solution may be prepared in water.</li> <li>• For in-line drench or dip applications, treat fruit for 15-30 seconds and allow fruit to drain.</li> <li>• Fruit coatings may be applied separately after treatments of this product.</li> </ul>
In-Line Aqueous or Fruit Coating Spray Application	Alternaria rot and surface mold Bitter rot Blue mold Bull's-eye rot Gray mold Phacidiopycnis rot Rhizopus rot Sphaeropsis rot White rot	16 fl oz/200,000 lb of fruit	<ul style="list-style-type: none"> <li>• Ensure proper coverage of the fruit.</li> <li>• Mix the fungicide solution in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated.</li> <li>• Use T-jet, CDA, or similar application system.</li> </ul>
<p><b>Restrictions:</b></p> <p>1) Do not make more than one post-harvest application of Academy to the fruit. Apply either once before storage or once after storage, just prior to marketing.</p> <p>2) Do not make more than two post-harvest applications of fludioxonil-containing products to the same fruit.</p> <ul style="list-style-type: none"> <li>• Ensure the Academy solution remains in suspension by using agitation.</li> </ul>			

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

### **Pesticide Storage**

Do not store near heat or open flame. Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

### **Pesticide Disposal**

Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

### **Container Handling (less than or equal to 5 gallons)**

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling (greater than 5 gallons)**

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling (greater than 5 gallons)**

**Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

Academy™, the ALLIANCE FRAME,  
the SYNGENTA Logo, and the PURPOSE ICON  
are trademarks of a Syngenta Group Company.



Viton™ is a trademark of The Chemours Company FC, LLC.

©2022 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1529A-L1A 0222**  
**4170570**



DIFENOCONAZOLE	GROUP	3	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE



## Fungicide

Active Ingredient:	
Difenoconazole:*	20.9%
Fludioxonil:**	12.5%
Other Ingredients:	66.6%
Total:	100.0%

\*CAS No. 119446-68-3

\*\*CAS No. 131341-86-1

Academy is a flowable suspension concentrate.

Academy contains 2.06 pounds of difenoconazole and 1.23 pounds of fludioxonil per gallon.

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1529

EPA Est. 100-NE-001

Academy™ and the Syngenta logo are trademarks of a Syngenta Group Company.

Viton™ is a trademark of The Chemours Company FC, LLC.

©2022 Syngenta

Manufactured for:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

SCP 1529A-L1A 0222  
4170570

**2.5 gallons**  
Net Contents

## KEEP OUT OF REACH OF CHILDREN. CAUTION

### FIRST AID

If **swallowed**: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**HOTLINE NUMBER**: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call **1-800-888-8372**.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

#### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, Viton™ ≥ 14 mils
- Shoes plus socks
- Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N\*, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N\*, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters

#### Environmental Hazards

This product is toxic to fish, mammals, aquatic invertebrates, oysters, and shrimp. Do not contaminate water when disposing of equipment washwater or rinsate. DO NOT allow contaminated waste water from the processing areas to enter lakes, streams, ponds or other waters. Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

### Physical or Chemical Hazards

Do not use or store near heat or open flame. Do not use or mix with oxidizing agents.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage**: Do not store near heat or open flame. Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

**Pesticide Disposal**: Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

**Container Handling**: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

**syngenta**®

