

PULL HERE TO OPEN ►

## RESTRICTED USE PESTICIDE DUE TO ACUTE INHALATION TOXICITY

FOR SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP 4A | 6 INSECTICIDES

 **Avicta® Duo COT202**

**syngenta®**

### Nematicide/Insecticide

A Seed Treatment Product to Protect Cotton Seedlings from Early Season Nematode and Insect Damage

For use only in Syngenta-certified cottonseed treatment facilities and only in seed treatment equipment with closed transfer and application systems. Not for use in hopper box, planter box, slurry box, or other farmer-applied applications.

*Active Ingredients:*

Abamectin (CAS No. 71751-41-2) .....	11.2%
Thiamethoxam (CAS No. 153719-23-4) .....	28.0%

<i>Other Ingredients:</i> .....	60.8%
---------------------------------	-------

<i>Total:</i> .....	100.0%
---------------------	--------

Avicta® Duo COT202 is a flowable concentrate containing 1.11 pounds of abamectin and 2.78 pounds of thiamethoxam per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

### WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1538 EPA Est. 100-NE-001

SCP 1538A-L1A 1016

4072958

Product ID

**55887**

**15 gallons**  
Net Contents

®

FIRST AID	
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<p style="text-align: center;"><b>HOT LINE NUMBER</b>  For 24-Hour Medical Emergency Assistance (Human or Animal)  Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)  Call  <b>1-800-888-8372</b></p>	

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### WARNING/AVISO

May be fatal if inhaled. Do not breathe vapor. May be fatal if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

This product must be used only in seed treatment equipment with closed transfer application systems.

The following personal protective equipment and clothing are required when using this product:

#### Personal Protective Equipment (PPE)

##### Seed Treatment Workers

**Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) must wear:**

- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils
- Protective eyewear
- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant apron
- Shoes and socks
- NIOSH-approved respirator with any N, R, or P filter with NIOSH-approval number prefix TC-84A; or a NIOSH-approved power air purifying respirator with HE filter with NIOSH-approval number prefix TC-21C

*continued...*

## PRECAUTIONARY STATEMENTS (*continued*)

### **Workers Filling Seed Containers/Baggers and Bag Sewers must wear:**

- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Protective eyewear
- Coveralls worn over long-sleeved shirt and long pants
- Shoes and socks
- NIOSH-approved respirator with any N, R, or P filter with NIOSH-approval number prefix TC-84A; or a NIOSH-approved power air purifying respirator with HE filter with NIOSH-approval number prefix TC-21C

### **Workers involved with the clean-up and maintenance of seed treatment equipment must wear:**

- Chemical-resistant gloves (minimum length 12 inches) made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils. Important: Tape gloves down to the sleeves of the coveralls.
- Protective eyewear
- Chemical-resistant, hooded coveralls worn over long-sleeved shirt and long pants
- Shoes and socks
- NIOSH-approved respirator with an organic vapor (OV) cartridge with a combination N, R, or P filter with NIOSH-approval number prefix TC-84A; **OR** NIOSH-approved gas mask with an organic vapor canister with NIOSH-approval number prefix TC-14G; **OR** a NIOSH-approved powered air purifying respirator with organic vapor (OV) cartridge and combination HE filter with NIOSH-approval number prefix TC-23C

### **Multiple Task Workers must wear:**

(Multiple task workers perform multiple tasks in one day such as mixing, product application, filling seed containers/ bagging, bag sewing, and clean-up)

- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils. Important: when performing seed treatment equipment clean-up procedures, gloves must be at least 12 inches long and taped down to the sleeves of the coveralls.
- Protective eyewear
- Long-sleeved shirt and long pants
- When performing seed treatment equipment clean-up procedures, chemical-resistant, hooded coveralls over long-sleeved shirt and long pants are required.
- Shoes and socks
- NIOSH-approved respirator with any N, R, or P filter with NIOSH-approval number prefix TC-84A; or a NIOSH-approved power air purifying respirator with HE filter with NIOSH-approval number prefix TC-21C

## **User Safety Requirements**

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### **User Safety Recommendations**

#### **Users should:**

- Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **Environmental Hazards**

This pesticide is toxic to fish and wildlife and highly toxic to aquatic invertebrates. Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading. Do not contaminate water when disposing of equipment washwater or rinsate.

#### **Pollinator Precautions**

Thiamethoxam is highly toxic to bees exposed to direct treatment, and effects may be possible as a result of exposure to translocated residues in blooming crops.

#### **Groundwater Advisory**

This product contains thiamethoxam which has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## DIRECTIONS FOR USE

### RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Restrictions:

- Avicta Duo COT202 is only for use in Syngenta-certified seed treatment facilities: Use in hopper box, planter box, slurry box, or other farmer-applied applications is not permitted.
- Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Conduct germination tests with a representative sample of the seed lot before committing the total seed lot to a selected seed treatment. Because seed quality and seed storage conditions are beyond the control of Syngenta, no claims are made to guarantee the germination of seed or propagating material for all crops.
- Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 180.1001. Refer to the section on this label **REQUIRED SEED CONTAINER STATEMENTS** for language that must appear on the tag that is attached to the container of treated seed.

#### PRODUCT INFORMATION

Avicta Duo COT202 is a cotton seed treatment nematicide/insecticide that contains the active ingredients abamectin and thiamethoxam. When Avicta Duo COT202 is applied to cotton seed, seedlings are protected from early season nematode damage and damage from the insect pests specified in the Crop Use Directions.

To add early season disease protection, the seed treatment fungicides specified in the Crop Use Directions are approved for use on cotton seed and may also be mixed with Avicta Duo COT202. Follow the most restrictive requirements and precautions for all products used in all mixtures. (See **Avicta Duo COT202 in Tank Mixtures**.)

#### REQUIRED SEED CONTAINER STATEMENTS

User is responsible for ensuring that the seed container meets all requirements under the Federal Seed Act.

The Federal Seed Act requires that the container of seed treated with Avicta Duo COT202 must be labeled with the following statements:

- This seed has been treated with abamectin nematicide and thiamethoxam insecticide.
- Do not use treated seed for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on the container of cottonseed treated with Avicta Duo COT202.

- Store away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and rubber gloves when handling treated seed.
- Groundwater Advisory: This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- **Pollinator Precautions:** Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.
- Plant Avicta Duo COT202-treated cottonseed based upon the local planting dates and soil temperatures that are recommended by your state agricultural extension agent.
- Treated seed must be planted into the soil at a depth greater than 1 inch.

- Treated seed exposed on soil surface may be hazardous to wildlife; cover or collect treated seed that are spilled during loading and in areas such as row ends.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging in accordance with local requirements.
- Do not use at a rate that will result in more than 0.075 lb thiamethoxam per acre per year (34.0 grams ai/A/yr) as a seed treatment application. This seed has been treated with thiamethoxam at 0.375 mg ai/seed, and 0.15 mg of abamectin per seed.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed, and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Allow a minimum of 45 days after planting Avicta Duo COT202 treated cottonseed before treating the cotton with a neonicotinoid insecticide.
- In the event of crop failure or after harvest of cotton grown from Avicta Duo COT202 treated cottonseed, the following crops may be replanted immediately:

Immediate Plant Back
<p>Alfalfa</p> <p>Cereal Grains: Barley, Buckwheat, Corn (field, pop, and sweet), Oat, Pearl Millet, Proso Millet, Rice (Dry-seeded), Rye, Sorghum, Teosinte, Triticale, Wheat, and Wild Rice</p> <p>Canola</p> <p>Cotton</p> <p>Cucurbit Vegetables Crop Group 9</p> <p>Fruiting Vegetables Crop Group 8</p> <p>Head and Stem <i>Brassica</i> Crop Subgroup 5A</p> <p>Leafy Brassica Greens Crop Subgroup 5B</p> <p>Leafy Vegetables (Except <i>Brassica</i> Vegetables) Crop Group 4</p> <p>Legume Vegetables (Succulent or Dried, Including Soybean) Crop Group 6</p> <p>Mint: Peppermint and Spearmint</p> <p>Oilseed Crops: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower</p> <p>Onion, Dry Bulb</p> <p>Peanut</p> <p>Potato</p> <p>Root Vegetables Crop Subgroup 1A</p> <p>Strawberry</p> <p>Sunflower</p> <p>Tobacco</p> <p>Tuberous and Corm Vegetables (Except Potato) Crop Subgroup 1D</p>

- For any other crop, the minimum plant back interval is 120 days from the date the Avicta Duo COT202 treated cotton seeds were planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

The following statement is required on containers of corn seed treated with Avicta Duo COT202:

- In areas with a history of high thrips pressure or for cotton grown in North Carolina, South Carolina, and Virginia, apply a foliar insecticide that is approved for thrips control, when cotton is between the 1st and 3rd leaf stage.

## RESISTANCE MANAGEMENT

GROUP	4A	6	INSECTICIDES
-------	----	---	--------------

Avicta Duo COT202 contains thiamethoxam, a Group 4A insecticide, and abamectin, a Group 6 insecticide. Thiamethoxam is a systemic insecticide belonging to the neonicotinoid class of chemistry which includes nicotinic acetylcholine receptor (nAChR) agonists. Abamectin belongs to the avermectin class of chemistry which affects nerves and muscles.

Insect populations may contain individuals naturally resistant to Group 4A or 6 insecticides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, use sound resistance management strategies established for the crop and use area.

Base seed treatment on an integrated pest management program that includes field sanitation, historical information related to pesticide use, careful selection of pest-tolerant crop varieties, scouting, and management practices which optimize populations of natural enemies of insect pests such as within-field refuge (untreated areas). Sound management programs also consider cultural and biological control practices.

### In order to maintain susceptibility to these classes of chemistry:

- Use products at their full specified doses.
- Use appropriate, well-maintained equipment. Use specified water volumes and apply at optimal temperatures in order to obtain optimal treatment.
- When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.
- Avoid using a single active ingredient or mode of action (same insecticide group) exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, use a treatment window approach. A treatment window is a period of time defined by the stage of crop development and the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, single or consecutive applications may be made using seed, in-furrow, or foliar treatments unless otherwise excluded by product labels. Do not exceed the maximum amount of this insecticide's mode of action allowed per growing season.
- Following a treatment window of this insecticide's mode of action, rotate to a treatment window of effective products with a different mode of action before making additional applications of this insecticide.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for the crop and use area.

Syngenta encourages responsible product stewardship to ensure effective long term control of the insect pests on this label.

### For additional information on Insect Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481
- Contact your local Cooperative Extension Service specialist, pest control advisor, or certified crop advisor.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org/>

## ROTATIONAL CROP RESTRICTIONS

In the event of crop failure or after harvest of cotton grown from Avicta Duo COT202 treated cottonseed, the following crops may be replanted immediately:

Immediate Plant Back
<p>Alfalfa</p> <p>Cereal Grains: Barley, Buckwheat, Corn (field, pop, and sweet), Oat, Pearl Millet, Proso Millet, Rice (Dry-seeded), Rye, Sorghum, Teosinte, Triticale, Wheat, and Wild Rice</p> <p>Canola</p> <p>Cotton</p> <p>Cucurbit Vegetables Crop Group 9</p> <p>Fruiting Vegetables Crop Group 8</p> <p>Head and Stem <i>Brassica</i> Crop Subgroup 5A</p> <p>Leafy Brassica Greens Crop Subgroup 5B</p> <p>Leafy Vegetables (Except <i>Brassica</i> Vegetables) Crop Group 4</p> <p>Legume Vegetables (Succulent or Dried, Including Soybean) Crop Group 6</p> <p>Mint: Peppermint and Spearmint</p> <p>Oilseed Crops: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower</p> <p>Onion, Dry Bulb</p> <p>Peanut</p> <p>Potato</p> <p>Root Vegetables Crop Subgroup 1A</p> <p>Strawberry</p> <p>Sunflower</p> <p>Tobacco</p> <p>Tuberous and Corm Vegetables (Except Potato) Crop Subgroup 1D</p>

For any other crop, the minimum plant back interval is 120 days from the date the Avicta Duo COT202 treated cotton seeds were planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

## MIXING AND APPLICATION INSTRUCTIONS

- Consult the Syngenta Seed Treatment representative or the manufacturer of the application equipment you plan to use regarding the suitability of this application and equipment calibration and operating instructions.
- Always treat seed in a well-ventilated area. Refer to the **Precautionary Statements** section of this label for personal protective equipment requirements and other safety precautions.
- Prepare no more mixture than is needed for the immediate operation.
- Avicta Duo COT202 must be thoroughly recirculated in its original container before using. (See **Mixing Instructions**.)
- Apply Avicta Duo COT202 as a water-based slurry utilizing standard slurry seed treatment equipment that will provide uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of nematode and insect control.

- Thoroughly mix the required amount of Avicta Duo COT202 and other seed treatment products desired with the amount of water required for the slurry treater and the dilution rate. (See **Avicta Duo COT202 in Tank Mixtures.**) Follow the most restrictive requirements and precautions for all products used in mixtures.
- Use an EPA-approved dye or colorant that imparts an unnatural color to the seed.
- Allow seed to dry before bagging /filling containers, and store treated seed away from feed and foodstuffs.

#### **Avicta Duo COT202 in Tank Mixtures:**

It is the pesticide user's responsibility to ensure that all products in a tank mix are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, plant back interval from another).

#### **Compatibility:**

Before mixing Avicta Duo COT202 with other seed treatment products, determine if all of the products are compatible by conducting a compatibility test:

1. In a clear glass container thoroughly mix all of the intended seed treatment products with the appropriate amount of water, and allow the mixture to sit for one hour.
2. Remix the solution and observe for incompatibility.
3. If compatibility is acceptable, follow the mixing instructions in the **Mixing Instructions** section on this label.

#### **Mixing Instructions:**

##### **A. Avicta Duo COT202 Alone**

1. To dispense Avicta Duo COT202 into a mix tank, first attach a pump with fittings (example: Micro Matic) that match the fittings on the Avicta Duo COT202 container.
2. Recirculate product. Recirculation time should be calculated based upon the ability of the pump to completely recirculate the volume of the tank 5-10 times within a 30-minute time period.
3. After the initial recirculation, a short recirculation period of 5 minutes per day is recommended.
4. Attach the dispensing hose to a dry-lock type of fitting that is permanently attached to the mix tank lid.
5. Add 1/4-1/2 of the required amount of water to the mix tank.
6. With the agitator running, add the Avicta Duo COT202 to the tank.
7. Continue agitation while adding the remainder of the water.
8. Begin application of the solution after the Avicta Duo COT202 has completely dispersed into the mix water.
9. Maintain agitation until all of the mixture has been applied.

##### **B. Avicta Duo COT202 - Tank Mixing Instructions:**

1. Add 1/2 of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred.
2. If using wettable powders and/or products in water-soluble packaging, add water-soluble packages first to clean water and allow them to completely disperse before adding Avicta Duo COT202 or other products.
3. Allow each tank mix partner to completely disperse before adding the next product.
4. Add the remaining amount of water and agitate.
5. Maintain agitation until the entire slurry mixture has been used.

## CROP USE DIRECTIONS

### Avicta Duo COT202 on Cottonseed

Avicta Duo COT202 is a seed treatment for the seed of delinted cotton. Avicta Duo COT202 will provide cotton seedlings early season protection from damage caused by nematodes and the insects specified in the following table. Tank mixing Avicta Duo COT202 with Dynasty® CST fungicide will add early season disease protection. Refer to the following table for application directions.

#### Application

Pests /Diseases	Amount of Product to Apply*	Instructions
<b>Nematodes and these insects:</b>  Cotton aphid Cotton fleahopper <sup>1</sup> Plant bugs <sup>1</sup> Tobacco thrips Western flower thrips Wireworm  <i><sup>1</sup>Avicta Duo COT202 only provides suppression of these pests. Suppression means either erratic control ranging from good to poor or consistent control at a level below that which is generally considered acceptable for commercial control.</i>	Apply Avicta Duo COT202 to cottonseed at a rate to achieve:  0.15 mg abamectin / 0.375 mg thiamethoxam per seed	To control early season nematodes and these insect pests apply the amount of Avicta Duo COT202 specified in the center column.  <b>NOTE:</b> <ul style="list-style-type: none"> <li>• *1 fl oz of Avicta Duo COT202 contains 3.93 grams of abamectin and 9.84 grams of thiamethoxam.</li> <li>• The typical density of Avicta Duo COT202 is 9.93 lb per gallon.</li> </ul>
<b>Disease Protection – Early Season</b>	Apply 3.1 – 3.95 fl oz/100 lb seed (0.025 – 0.032 mg ai/ seed) of Dynasty CST.	To add early season disease control in cotton, cottonseed may be treated with a mixture of Avicta Duo COT202 plus Dynasty CST. Follow the most restrictive requirements and precautions for all products used in mixtures.

## CLEANING SEED TREATMENT EQUIPMENT

Do not use compressed air hoses to remove the dust from the seed treating equipment or the area around the equipment; use a central vacuum system or a HEPA filter portable vacuum cleaner.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### Pesticide Storage

Store in a cool, dry place.

#### Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

#### Container Handling

**Nonrefillable container. Do not reuse or refill this container.** Syngenta's authorized agent will pick up the empty container and handle its disposal. Contact Syngenta or its authorized agent to make arrangements for this service and store empty container in a secure area until pick up.

Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then contact Syngenta or its authorized agent for disposal.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean-up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

Avicta®, Dynasty®, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company



Viton® is a trademark of E. I. duPont de Nemours & Co., Inc.

Protected by U.S. Patent Nos. 5,852,012, & 6,875,727.

©2016 Syngenta

For non-emergency (e.g. current product information) call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1538A-L1A 1016  
4072958**

# RESTRICTED USE PESTICIDE DUE TO ACUTE INHALATION TOXICITY

FOR SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP 4A 6 INSECTICIDES



## Nematicide/ Insecticide

A Seed Treatment Product to Protect  
Cotton Seedlings from Early Season  
Nematode and Insect Damage

For use only in Syngenta-certified cottonseed  
treatment facilities and only in seed treatment  
equipment with closed transfer and application  
systems. Not for use in hopper box, planter box,  
slurry box, or other farmer-applied applications.

### Active Ingredients:

Abamectin (CAS No. 71751-41-2) ..... 11.2%  
Thiamethoxam (CAS No. 153719-23-4) ..... 28.0%

Other Ingredients: ..... 60.8%  
Total: ..... 100.0%

Avicta® Duo COT202 is a flowable concentrate  
containing 1.11 pounds of abamectin and 2.78  
pounds of thiamethoxam per gallon.

See additional precautionary statements and  
directions for use in booklet.

EPA Reg. No. 100-1538

EPA Est. 100-NE-001

Product ID

# 55887

Avicta® and the Syngenta Logo are Trademarks  
of a Syngenta Group Company

Protected by U.S. Patent Nos. 5,852,012, &  
6,875,727.

©2016 Syngenta

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

SCP 1538A-L1A 1016  
4072958

## 15 gallons

Net Contents

KEEP OUT OF REACH  
OF CHILDREN.

## WARNING/ AVISO

Si usted no entiende la etiqueta, busque a  
alguien para que se la explique a usted en  
detalle. (If you do not understand the label,  
find someone to explain it to you in detail.)

### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

#### WARNING/AVISO

May be fatal if inhaled. Do not breathe vapor.  
May be fatal if swallowed. Avoid contact with  
skin, eyes, or clothing. Wash thoroughly with  
soap and water after handling and before  
eating, drinking, chewing gum, using tobacco  
or using the toilet. Remove and wash contam-  
inated clothing before reuse.

This product must be used only in seed treat-  
ment equipment with closed transfer applica-  
tion systems.

#### FIRST AID

If inhaled: Move person to fresh air. If person  
is not breathing, call 911 or an ambulance,  
then give artificial respiration, preferably  
mouth-to-mouth if possible. Call a poison  
control center or doctor for further treat-  
ment advice. If swallowed: Call poison con-  
trol center or doctor immediately for treat-  
ment advice. Have person sip a glass of water  
if able to swallow. Do not induce vomiting  
unless told to do so by the poison control  
center or doctor. Do not give anything by  
mouth to an unconscious person.

Have the product container or label with  
you when calling a poison control center or  
doctor, or going for treatment.

**HOT LINE NUMBER:** For 24-Hour Medical  
Emergency Assistance (Human or Animal) Or  
Chemical Emergency Assistance (Spill, Leak,  
Fire or Accident) Call 1-800-888-8372.

**Environmental Hazards:** This pesticide is toxic  
to fish and wildlife and highly toxic to aquatic  
invertebrates. Treated seeds exposed on the  
soil surface may be hazardous to wildlife.  
Cover or collect seeds spilled during loading.  
Do not contaminate water when disposing of  
equipment washwater or rinsate.

**Pollinator Precautions:** Thiamethoxam is highly  
toxic to bees exposed to direct treatment, and  
effects may be possible as a result of exposure  
to translocated residues in blooming crops.

**Groundwater Advisory:** This product contains  
thiamethoxam which has properties and char-  
acteristics associated with chemicals detected

in groundwater. This chemical may leach into  
groundwater if used in areas where soils are  
permeable, particularly where the water  
table is shallow.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed  
by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place.

**Pesticide Disposal:** Pesticide wastes are  
acutely hazardous. Improper disposal of  
excess pesticide, spray mixture or rinsate is  
a violation of Federal Law. If these wastes  
cannot be disposed of by use according to  
label instructions, contact your State Pesti-  
cide or Environmental Control Agency, or  
the Hazardous Waste Representative at the  
nearest EPA Regional Office for guidance.

#### Container Handling

**Nonrefillable container.** Do not reuse or  
refill this container. Syngenta's authorized  
agent will pick up the empty container  
and handle its disposal. Contact Syngenta  
or its authorized agent to make arrange-  
ments for this service and store empty  
container in a secure area until pick up.

Triple rinse container promptly after  
emptying. Triple rinse as follows: Empty  
the remaining contents into application  
equipment or a mix tank. Fill the container  
1/4 full with water. Replace and tighten  
closures. Tip container on its side and roll  
it back and forth, ensuring at least one  
complete revolution, for 30 seconds. Stand  
the container on its end and tip it back  
and forth several times. Turn the container  
over onto its other end and tip it back  
and forth several times. Empty the rinsate  
into application equipment or a mix tank  
or store rinsate for later use or disposal.  
Repeat this procedure two more times.  
Then contact Syngenta or its authorized  
agent for disposal.

For minor spills, leaks, etc., follow all pre-  
cautions indicated on this label and clean-  
up immediately. Take special care to avoid  
contamination of equipment and facilities  
during clean-up procedures and disposal  
of wastes.

**CONTAINER IS NOT SAFE FOR FOOD, FEED,  
OR DRINKING WATER!**

