

MEFENOXAM GROUP 4 FUNGICIDE

PULL HERE TO OPEN ►



Orondis® Gold B

syngenta®

Fungicide

For the control of certain diseases in listed crops caused by the Oomycete class of fungi

Active Ingredient:

Mefenoxam* 45.3%

Other Ingredients: 54.7%

Total: 100.0%

*CAS No. 70630-17-0 and 69516-34-3

Orondis® Gold B is formulated as a soluble concentrate and contains 4 lb active ingredient per gallon.

SCP 1202B-L1H 1120 4134440

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1202 EPA Est. 072344-MO-004

Product of Switzerland

2.5 quarts
(80 fluid ounces)

Net Contents



TABLE OF CONTENTS

- 1.0 FIRST AID**
- 2.0 PRECAUTIONARY STATEMENTS**
 - 2.1 Hazards to Humans and Domestic Animals**
 - 2.2 Personal Protective Equipment (PPE)**
 - 2.2.1 User Safety Requirements
 - 2.2.2 Engineering Controls
 - 2.2.3 User Safety Recommendations
 - 2.3 Environmental Hazards**
 - 2.3.1 Groundwater Advisory
 - 2.3.2 Surface Water Advisory
 - 2.4 Physical or Chemical Hazards**
- DIRECTIONS FOR USE**
- 3.0 PRODUCT INFORMATION**
 - 3.1 Integrated Pest Management (IPM)**
 - 3.2 Resistance Management**
- 4.0 APPLICATION DIRECTIONS**
 - 4.1 Methods of Application**
 - 4.1.1 Band Application
 - 4.1.2 In-Furrow Application

DIRECTIONS FOR USE

- 4.2 Application Equipment**
 - 4.2.1 Nozzles
 - 4.2.2 Pump
- 4.3 Application Volume and Spray Coverage**
- 4.4 Mixing Directions**
 - 4.4.1 Orondis Gold B Alone
 - 4.4.2 Tank-Mix Precautions
 - 4.4.3 Tank-Mix Compatibility Test
 - 4.4.4 Orondis Gold B in Tank Mixtures
- 4.5 Application through Irrigation Systems (Chemigation)**
 - 4.5.1 Chemigation Restrictions
 - 4.5.2 Operating Instructions for Public Water Systems
 - 4.5.3 Specific Instructions for Public Water Systems
 - 4.5.4 Application Directions for Irrigation Systems

continued...

DIRECTIONS FOR USE *(continued)*

5.0 REPLANT AND ROTATIONAL CROP RESTRICTIONS

5.1 Replanting

5.2 Rotational Crop Restrictions

6.0 RESTRICTIONS AND PRECAUTIONS

6.1 Use Restrictions

6.2 Use Precautions

6.3 Spray Drift Management

6.4 Drift Reduction Advisory Information

6.4.1 Importance of Droplet Size

6.4.2 Controlling Droplet Size – Ground Boom

6.4.3 Controlling Droplet Size – Aircraft

6.4.4 Boom Height – Ground Boom

6.4.5 Release Height - Aircraft

6.4.6 Shielded Sprayers

6.4.7 Temperature and Humidity

6.4.8 Temperature Inversions

6.4.9 Wind

7.0 CROP USE DIRECTIONS FOR SOIL INJECTED OR SOIL INCORPORATED APPLICATIONS

7.1 Berry and Small Fruit

7.1.1 Bushberry, Crop Subgroup 13-07B

7.1.2 Caneberry, Crop Subgroup 13-07A

7.2 Cucurbit Vegetables, Crop Group 9

7.3 Fruiting Vegetables

7.3.1 Crop Group 8, except Tomato

7.3.2 Tomato

7.4 Leafy Vegetables (except Brassica), Crop Group 4

7.5 Root and Tuber Vegetables

7.5.1 Crop Group 1, except Carrot, Ginseng, Potato and Sugar Beet

7.5.2 Ginseng

7.5.3 Potato

7.6 Tobacco

8.0 CROP USE DIRECTIONS FOR SOIL DIRECTED APPLICATIONS

8.1 Asparagus

8.2 Berry and Small Fruit

8.2.1 Bushberry, Crop Subgroup 13-07B

8.2.2 Caneberry, Crop Subgroup 13-07A

8.3 Cucurbit Vegetables, Crop Group 9

8.4 Fruiting Vegetables

8.4.1 Crop Group 8, except Tomato

8.4.2 Tomato

8.5 Root and Tuber Vegetables

8.5.1 Potato

8.6 Tobacco

9.0 STORAGE AND DISPOSAL

10.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

1.0 FIRST AID

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by the poison control center or doctor.• Do not give anything to an unconscious person.

FIRST AID (continued)	
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOTLINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-888-8372	

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton® \geq 14 mils
- Shoes plus socks

2.2.1 USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.2.2 ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

2.2.3 USER SAFETY RECOMMENDATIONS

User Safety Recommendations

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.3 Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

2.3.1 GROUNDWATER ADVISORY

Mefenoxam is known to leach through the soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

2.3.2 SURFACE WATER ADVISORY

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

2.4 Physical or Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT APPLY this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS). **Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.** For crop uses that fall under this REI, see **Section 8.0, Soil-Directed Applications**. **Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area immediately if there will be no contact with anything that has been treated. For crop uses that fall under this Exception, see **Section 7.0, Soil-Injected or Soil-Incorporated Applications**. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: • Coveralls • Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils • Shoes plus socks

3.0 PRODUCT INFORMATION

Orondis Gold B is a systemic fungicide for use on selected crops to control certain diseases caused by members of the Oomycete class of fungi.

To ensure maximum activity on soilborne pathogens, Orondis Gold B must be moved into the seed or root zone of the plant. Placement in the seed or root zone includes in-furrow sprays, soil injections, crown dips. Incorporation includes preplant incorporated applications, soil drenches, or shank applications.

For soil surface sprays, rainfall will move the fungicide into the seed or root zone, but if rain is not expected within 24 hours after application, mechanically incorporate (before planting) or sprinkler irrigate (after planting) with 1/2 to 1 inch of water.

Under conditions conducive to severe disease pressure, additional fungicide applications may be applied using an alternate fungicide registered for the crop/disease appearing on this label.

Where rate ranges are specified on this label, use the higher specified rate when heavy disease pressure is expected and the lower specified rate when disease pressure is expected to be light, unless otherwise noted.

EFFICACY

Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of Orondis Gold B has been used. If resistant isolates to Group 4 fungicides are present, efficacy can be reduced. Under high disease pressure, it is recommended to use the highest specified rate and shortest specified interval when needed.

CROP TOLERANCE

Plant tolerance has been found acceptable for all crops on the label, however, not all possible tank-mix combinations have been tested under all conditions. When possible, it is recommended to test the combinations on a small portion of the crop to ensure a phytotoxic response will not occur as a result of application.

3.1 Integrated Pest Management (IPM)

Orondis Gold B should be integrated into an overall disease and pest management strategy (IPM) whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area.

3.2 Resistance Management

MEFENOXAM	GROUP	4	FUNGICIDE
-----------	-------	---	-----------

For resistance management, Orondis Gold B contains a Group 4 fungicide. Any fungal population may contain individuals naturally resistant to Orondis

Gold B and other Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Orondis Gold B or other Group 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crop and pathogens.

- For further information or to report suspected resistance contact Syngenta at 1-866-Syngent(a) (866-796-4368). You can also contact your pesticide distributor or university extension specialist to report resistance.

4.0 APPLICATION DIRECTIONS

4.1 Methods of Application

Applications with Orondis Gold B are permitted by ground, by air and via chemigation as specified in **Sections 7.0 and 8.0** unless otherwise restricted in **Section 6.1**. Ground application includes broadcast sprays, soil incorporation, banded and in-furrow applications as well as soil injections and crown dips. Incorporation includes preplant incorporated, soil drenches or shank applications.

For band applications refer to **Section 4.1.1** to calculate the amount of Orondis Gold B and water volume needed. For In-furrow applications, refer to **Section 4.1.2** for the amount of product to use with common row spacings. Refer to **Section 4.5** for details of application by chemigation.

4.1.1 BAND APPLICATION

Application rates in the **Sections 7.0 and 8.0** are generally expressed as an amount per acre which refers to the total crop area to be treated. If using a banded application, use proportionally less product using the formula below:

$$\frac{\text{band width in inches}}{\text{row spacings in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

4.1.2 IN-FURROW APPLICATION

The following table provides common row spacing and the amount of Orondis Gold B to apply per acre.

Use Rate fl oz/1,000 row feet (oz ai/1,000 row feet)	Orondis Gold B fl oz/A								
	20-inch rows	22-inch rows	24-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.08 (0.038)	2.1	1.9	1.7	1.4	1.3	1.2	1.1	1.1	1.0
0.15 (0.078)	3.9	3.6	3.2	2.6	2.4	2.3	2.2	2.0	1.9
0.28 (0.14)	7.3	6.7	6.1	4.8	4.6	4.3	4.0	3.8	3.7
0.42 (0.21)	11.0	10.0	9.1	7.3	6.8	6.5	6.1	5.8	5.4

20" = 26,136 row ft/A, 22" = 23,760 row ft/A, 24" = 21,780 row ft/A, 30" = 17,424 row ft/A, 32" = 16,315 row ft/A, 34" = 15,374 row ft/A, 36" = 14,520 row ft/A, 38" = 13,754 row ft/A, 40" = 13,068 row ft/A

4.2 Application Equipment

- Spray equipment configuration should be arranged to provide accurate application, and minimize potential for spray drift.
- To ensure accuracy, calibrate sprayer before each use.
- For information on spray equipment and calibration, consult spray equipment manufacturers and/or state recommendations.
- All ground/aerial/chemigation application equipment must be properly maintained and calibrated using appropriate carriers.

4.2.1 NOZZLES

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump should be 16-mesh or coarser.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

4.2.2 PUMP

- Use a pump with capacity to:
 - o Maintain 35-40 psi at nozzles
 - o Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.

- Use a jet agitator or liquid sparger tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

4.3 Application Volume and Spray Coverage

- Apply by ground or air in sufficient water or liquid fertilizer to provide uniform coverage of the soil surface.
- For maximum effectiveness, Orondis Gold B must be moved into the seed or root zone of the plant.
- For in-furrow application, apply as an in-furrow spray in 3-7 gallons per acre of water at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered.
- For ground broadcast application, apply in a minimum of 20 gallons of water per acre, unless specified otherwise. If rain is not expected within 24 hours after application, mechanically incorporate (before planting) or sprinkler irrigate (after planting) with $\frac{1}{2}$ to 1 inch of water.
- For aerial application, apply in a minimum of 3 gallons of water per acre, unless specified otherwise. Avoid application when uniform coverage cannot be obtained or when excessive spray drift may occur.

4.4 Mixing Directions

1. Thoroughly clean spray equipment before using this product.
2. Prepare no more spray mixture than is needed for the immediate operation.
3. Keep product container tightly closed when not in use.
4. Agitate the spray solution before and during application.
5. Do not let the spray mixture stand overnight in the spray tank.
6. Rinse the spray equipment thoroughly after each day's use and dispose of pesticide rinsate by application to an already treated area.

4.4.1 ORONDIS GOLD B ALONE

1. Add 1/4 - 1/2 of the required amount of water to the spray or mixing tank.
2. With the agitator running add Orondis Gold B to the tank.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the spray solution after Orondis Gold B has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been sprayed.

4.4.2 TANK-MIX PRECAUTIONS

- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Tank mixes of Orondis Gold B with other pesticides, fertilizers, or any other additives not specifically labelled for use with Orondis Gold B

may result in tank mix incompatibility or unsatisfactory performance. In such cases, always check tank mix compatibility by conducting a jar test according to guidance in **Section 4.4.3** before actual tank mixing.

4.4.3 TANK-MIX COMPATIBILITY TEST

- Conduct a jar test using a 1 pt to 1 qt container with lid by adding water or other intended carrier such as a liquid fertilizer to the jar.
- Next, add the appropriate amount of pesticide(s) or tank mix partner(s) in their relative proportions based on recommended label rates. Add tank mix components separately in the order described in the tank-mixing section, **Section 4.4.4**. After each addition, shake or stir gently to thoroughly mix.
- After all ingredients have been added, put the lid on the jar, tighten and invert the jar 10 times to mix.
- After mixing, let the mixture stand 15 - 30 minutes and then examine for signs of incompatibility such as obvious separation, large flakes, precipitates, gels or heavy oily film on the jar.
- If the mixture remains mixed or can be remixed readily, it is physically compatible and can be used.
- If the mixture is incompatible, repeat the test using a compatibility agent at the recommended rate. Or, if applicable, slurry dry formulations in water before adding to the jar. If incompatibility is still observed after following these procedures, do not use the mixture.
- After compatibility testing is complete, dispose of any pesticide wastes in accordance with the storage and disposal section, **Section 9.0** of this label.

4.4.4 ORONDIS GOLD B IN TANK MIXTURES

1. Add 1/4 - 1/2 of the required amount of water to the spray or mixing tank.
2. With the agitator running, add the tank-mix partner(s) into the tank in the following order:
 - a. wettable powders
 - b. water dispersible granular products
 - c. liquid flowables
 - d. emulsifiable concentrates
3. Allow the material to completely dissolve and disperse into the mix water.
4. Continue agitation while adding the remainder of the water and Orondis Gold B to the spray tank.
5. Allow Orondis Gold B to completely disperse.
6. Spray the mixture with the agitator running.
7. Follow the precautions and limitations of the most restricted product in the tank mixture.

4.5 Application through Irrigation Systems (Chemigation)

4.5.1 CHEMIGATION RESTRICTIONS

- Use only on crops for which chemigation is specified on this label.
- Apply Orondis Gold B only through center pivot, solid set, hand move, moving wheel, micro-sprinkler, or drip irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Orondis Gold B must be applied on the schedule specified in the specific crop use directions, not according to the irrigation schedule.

Note: Orondis Gold B can affect many seal materials and should not be used at full strength. Leather seals are best, EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton, Buna-N, Neoprene, or PVC seals.

4.5.2 OPERATING INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

continued...

4.5.2 OPERATING INSTRUCTIONS FOR PUBLIC WATER SYSTEMS (continued)

3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

4.5.3 SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the

functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering device, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

4.5.4 APPLICATION DIRECTIONS FOR IRRIGATION SYSTEMS

- Apply Orondis Gold B only through center pivot, solid set, hand move, moving wheel, micro-sprinkler, or drip irrigation systems. Do not apply this product through any other type of irrigation system.
- Do not inject Orondis Gold B at full strength or deterioration of seals may occur. Use a dilution ratio of at least 15 parts water to 1 part Orondis Gold B in the tank mix.
- The chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.
- Good agitation should be maintained during the entire application period.
- Uniform coverage is required for good control.

5.0 REPLANT AND ROTATIONAL CROP RESTRICTIONS

5.1 Replanting

If replanting is necessary, additional applications of Orondis Gold B may be made, provided that the total amount of active ingredient in Orondis Gold B applied does not exceed the maximum allowed for the specific crop.

5.2 Rotational Crop Restrictions

The following crops may be planted at the specified interval following application of Orondis Gold B.

Crop, Crop Group or Crop Subgroup	Replant/Plant-Back Interval
Alfalfa (birdsfoot trefoil) Apple Artichoke, globe Asparagus Avocado Brassica (Cole) leafy vegetables (e.g. broccoli, cabbage, cauliflower) Bushberry, Subgroup 13-07B (e.g., blueberries, cranberries) Cacao Caneberry, Subgroup 13-07A (e.g. blackberry and raspberry) Citrus Clover Corn Cotton Cucurbit vegetables (e.g. cucumber, melons, squash)	0 days

continued...

5.2 Rotational Crop Restrictions (continued)

Crop, Crop Group or Crop Subgroup	Replant/Plant-Back Interval
Fruiting vegetables (e.g. tomato, peppers, eggplant) Ginseng Grapes Grass, forage, fodder and hay Herbs (fresh and dried) Hops Leafy vegetables, except Brassica (e.g. lettuce, spinach, celery) Legume vegetables (e.g. beans and peas, succulent and dried) Onions (dry bulb, garlic, and green) Peanuts Pineapples Root and tuber vegetables (e.g. potato, carrots, sugar beets) Soybeans Stone fruits Strawberries Sunflower Tobacco	0 days

Crop, Crop Group or Crop Subgroup	Replant/Plant-Back Interval
Tomato Tree nuts, Crop Group 14-12 Tropical fruit (e.g. papaya, mango) Wasabi	0 days
Cereal grains (except corn)	14 days
Crops not intended for food or feed	0 days
All other crops intended for food and feed	365 days

6.0 RESTRICTIONS AND PRECAUTIONS

See **Sections 7.0** and **8.0** for crop-specific Restrictions and Precautions.

6.1 Use Restrictions

- **DO NOT** use in greenhouses or other structures such as lath houses, float houses, and hydroponic facilities, unless specified on this label.
- **DO NOT** use for disease control in bedding plants, transplant trays, or nurseries except where specifically allowed in certain crop sections.
- **DO NOT** use as a foliar application unless specified on this label.
- **DO NOT** dip plants or roots, spray bare roots, or use a transplant water treatment with solutions containing Orondis Gold B except where specifically allowed in certain crop sections.

- Maximum usage when applying both mefenoxam- and metalaxyl-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.
- Soil-applied maximum application rates include all uses in **Section 7.0** and all uses in **Section 8.0** except foliar applications.

6.2 Use Precautions

- Avoid spray overlap as crop injury may occur.

6.3 Spray Drift Management

- AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.
- To avoid spray drift, do not apply when conditions favor drift beyond the target area.
- The interaction of many equipment and weather related factors determine the potential for spray drift.
- More information on managing spray drift can be found on the Syngenta Crop Protection website under Stewardship.

<http://www.syngentacropprotection-us.com/enviro/driftmanagement/>

6.4 Drift Reduction Advisory Information

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

6.4.1 IMPORTANCE OF DROPLET SIZE

- An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control.
- While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

6.4.2 CONTROLLING DROPLET SIZE – GROUND BOOM

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

6.4.3 CONTROLLING DROPLET SIZE – AIRCRAFT

- **Adjust Nozzles** - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

6.4.4 BOOM HEIGHT - GROUND BOOM

- Use the lowest boom height that is compatible with the spray nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

6.4.5 RELEASE HEIGHT - AIRCRAFT

- Higher release heights increase the potential for spray drift. When applying aerially to crops, do not spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

6.4.6 SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers.
- Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

6.4.7 TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

6.4.8 TEMPERATURE INVERSIONS

- Drift potential is high during a temperature inversion.
- Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind.
- The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator.
- Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.
- Avoid applications during temperature inversions.

6.4.9 WIND

- Drift potential generally increases with wind speed.
- AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

7.0 CROP USE DIRECTIONS FOR SOIL INJECTED OR SOIL INCORPORATED APPLICATIONS

- The restricted-entry interval (REI) for soil-injected or soil incorporated applications is 0 hours.

7.1 Berry and Small Fruit

7.1.1 BUSHBERRY, CROP SUBGROUP 13-07B

Crops (including all cultivars, varieties, and/or hybrids of these)			
<div> <div> Aronia berry Blueberry, highbush Blueberry, lowbush Buffalo currant Chilean guava Currant, black Currant, red </div> <div> Elderberry European barberry Gooseberry Highbush cranberry Honeysuckle, edible Huckleberry </div> <div> Jostaberry Juneberry (Saskatoon Berry) Lingonberry Native currant Salal Sea buckthorn </div> </div>			
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Root rot (<i>Phytophthora</i> spp.)	3.6 (1.8)	New Plantings: Apply after initial at-planting application.	Soil Application (band or via drip/micro-sprinkler irrigation) New Plantings: Reapply once during a period favorable for root rot. Use Orondis Gold B in conjunction with good cultural practices to minimize disease.
Resistance Management: • Refer to Section 3.2.			
Precaution: • Orondis Gold B will not revitalize plants showing moderate to severe root rot symptoms.			

continued...

7.1.1 BUSHBERRY, CROP SUBGROUP 13-07B (continued)

USE RESTRICTIONS	
1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: 3.6 pt/A (equivalent to 1.8 lb ai/A mefenoxam) 3) Minimum Application Interval: 90 days 4) Maximum Annual Application Rate: 7.2 pt/A/year (equivalent to 3.6 lb ai/A/year mefenoxam) a) DO NOT exceed 3.6 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products. 5) Pre-harvest Interval (PHI): 0 days	

7.1.2 CANEBERRY, CROP SUBGROUP 13-07A

Crops (including all cultivars, varieties, and/or hybrids of these)			
Blackberry Loganberry		Raspberry, black Raspberry, red	Raspberry, wild
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Root rot (<i>Phytophthora</i> spp.)	3.6 (1.8)	New Plantings: After initial at-planting application (Section 7.0).	Soil Application (band or via drip/micro-sprinkler irrigation) New Plantings: Reapply once during a period favorable for root rot. Use Orondis Gold B in conjunction with good cultural practices to minimize disease.
See directions for Established Plantings in 8.2.2 for subsequent applications.			

Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2.
Precaution: <ul style="list-style-type: none"> Orondis Gold B will not revitalize plants showing moderate to severe root rot symptoms.
USE RESTRICTIONS
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: 3.6 pt/A (equivalent to 1.8 lb ai/A mefenoxam) Minimum Application Interval: 90 days Maximum Annual Application Rate: 7.2 pt/A/year (equivalent to 3.6 lb ai/A/year mefenoxam) <ol style="list-style-type: none"> DO NOT exceed 3.6 lb ai/A/year of soil-applied and 0.2 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products. DO NOT use an adjuvant. Pre-harvest Interval (PHI): 45 days

7.2 Cucurbit Vegetables, Crop Group 9

Crops (including all cultivars, varieties, and/or hybrids of these)		
Chayote (fruit)	Muskmelon	Squash, summer
Chinese waxgourd (Chinese preserving melon)	Cantaloupe	Crookneck squash
Citron melon	Casaba	Scallop squash
Cucumber	Crenshaw melon	Straightneck squash
Gherkin	Golden pershaw melon	Vegetable marrow
Gourd, edible	Honeydew melon	Zucchini
Hyotan	Honey balls	Squash, winter
Cucuzza	Mango melon	Acorn squash
Hechima	Persian melon	Butternut squash
Chinese okra	Pineapple melon	Calabaza
Momordica spp.	Santa Claus melon	Hubbard squash
Balsam apple	Snake melon	Spaghetti squash
Balsam pear	True cantaloupe	Watermelon
Bittermelon	Pumpkin	
Chinese cucumber		

Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Damping off (<i>Pythium</i> spp.) Suppression: Phytophthora blight (<i>Phytophthora capsici</i>)	1.0 – 2.0 (0.5-1.0)	Preplant incorporated: At planting	For preplant incorporation , apply as a broadcast or banded application in water or liquid fertilizer and incorporate in the top 2 inches of soil. For soil spray applications , apply as a broadcast or banded application in water or liquid fertilizer at planting. For injection (drip irrigation) , inject Orondis Gold B into the irrigation water at the labeled rates. For banded applications, use a 7-inch band.
For soil directed applications, refer to Section 8.3			
Resistance Management: • Refer to Section 3.2 .			

continued...

7.2 Cucurbit Vegetables, Crop Group 9 (continued)

USE RESTRICTIONS	
1) Refer to Section 6.1 for additional product use restrictions.	
2) Maximum Single Application Rate: 2.0 pt/A (equivalent to 1.0 lb ai/A mefenoxam)	
3) Minimum Application Interval: 14 days	
4) Maximum Annual Application Rate: Soil: 2.0 pt/A/year (equivalent to 1.0 lb ai/A/year mefenoxam)	
a) DO NOT exceed 1.0 lb ai/A/year of soil-applied and 0.5 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.	
5) Pre-harvest Interval (PHI): 5 days	

7.3 Fruiting Vegetables

7.3.1 CROP GROUP 8, EXCEPT TOMATO

Crops (including all cultivars, varieties, and/or hybrids of these)		
Eggplant	Pepper	Pepper
Groundcherry	Bell	Pimento
Pepino	Chili	Sweet
	Cooking	Tomatillo

Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Damping off (<i>Pythium</i> spp.) Suppression: Crown rot (<i>Phytophthora capsici</i>)	1.0 (0.5)	Preplant or at-planting For direct seeded peppers , apply preplant or prior to emergence. For application by drip irrigation , make up to two additional applications on a 30-day schedule following initial application at planting. To control crown rot , apply before the plants are infected to obtain satisfactory control.	For soil spray applications , apply as a broadcast or banded application in water or liquid fertilizer preplant or at-planting. For banded applications, use a 12- to 16-inch band. For Injection (drip irrigation) , inject Orondis Gold B into the irrigation water.
For soil directed applications, refer to Section 8.4.1			
Resistance Management: • Refer to Section 3.2 .			
Precaution: • Plants already infected with <i>Phytophthora capsici</i> cannot be cured with Orondis Gold B.			

continued...

7.3.1 CROP GROUP 8, EXCEPT TOMATO (continued)

USE RESTRICTIONS	
1)	Refer to Section 6.1 for additional product use restrictions.
2)	Maximum Single Application Rate: 1.0 pt/A (equivalent to 0.5 lb/A mefenoxam)
3)	Minimum Application Interval: 30 days
4)	Maximum Annual Application Rate: 3.0 pt/A/year (equivalent to 1.5 lb ai/A/year mefenoxam) a) DO NOT exceed 1.5 lb ai/A/year of soil-applied and 0.5 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.
5)	Pre-harvest Interval (PHI): 7 days

7.3.2 TOMATO

Crops (including all cultivars, varieties, and/or hybrids)			
Tomato			
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Damping off (<i>Pythium</i> spp.)	1.0 – 2.0 (0.5-1.0)	At planting	Soil spray (broadcast or band): Apply in water or liquid fertilizer. For banded applications, use a 7-inch band.

Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Fruit rot Root rot (<i>Phytophthora</i> spp.) (<i>Pythium</i> spp.)	1.0 (0.5)	4-6 weeks after planting Apply a second drip irrigation as needed up to 1 week before harvest.	Injection (drip irrigation): Initiate control of fruit and root rot with a soil application as described above. Make subsequent applications by drip application according to the application timing schedule. For injected applications, base rate calculations on a 7-inch band.
For soil directed applications, refer to Section 8.4.2			
Resistance Management:			
• Refer to Section 3.2.			
USE RESTRICTIONS			
1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: 2.0 pt/A (equivalent to 1.0 lb ai/A mefenoxam) 3) Minimum Application Interval: 28 days 4) Maximum Annual Application Rate: 3.0 pt/A/year (equivalent to 1.5 lb ai/A/year mefenoxam) a) DO NOT exceed 1.5 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products and 0.5 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products. 5) Pre-harvest Interval (PHI): 7 days			

7.4 Leafy Vegetables (except Brassica), Crop Group 4

Crops (including all cultivars, varieties, and/or hybrids of these)		
Amaranth, Chinese spinach	Corn salad	Orach
Amaranth, leafy	Cress, garden	Parsley
Amaranth, tampala	Cress, upland	Purslane, garden
Arugula (Rocket)	Dandelion	Purslane, winter
Cardoon	Dock (sorrel)	Radicchio (red chicory)
Celery	Dillweed	Rhubarb
Celery, Chinese	Dock	Spinach
Celtuce	Endive (escarole)	Spinach, New Zealand
Chervil	Fennel, Florence (finocchio)	Spinach, vine
Chrysanthemum, edible-leaved	Lettuce, head and leaf	Swiss chard
Chrysanthemum, garland		

Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Damping off (<i>Pythium</i> spp.)	1.0 – 2.0 (0.5-1.0)	Preplant incorporated At planting	Preplant incorporated (broadcast or band): Apply in water or liquid fertilizer and mechanically incorporate in the top 2 inches of soil. Soil spray (broadcast or band): Apply in water or liquid fertilizer at planting. For banded applications, use a 7-inch band.
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2. 			

continued...

7.4 Leafy Vegetables (except Brassica), Crop Group 4 (continued)

USE RESTRICTIONS	
1)	Refer to Section 6.1 for additional product use restrictions.
2)	Maximum Single Application Rate: 2.0 pt/A (equivalent to 1.0 lb ai/A mefenoxam)
3)	Minimum Application Interval: 21 days - spinach only.
4)	Maximum Annual Application Rate: Soil: 2.0 pt/A/year (equivalent to 1.0 lb ai/A/year mefenoxam) a) For lettuce, DO NOT exceed 1.0 lb ai/A/year of soil-applied and 0.4 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products. b) For spinach, EITHER, DO NOT exceed 1.0 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products at planting and 0.25 lb ai/A/year of soil-applied post planting, shanked-in applications of mefenoxam- and metalaxyl-containing products. OR, DO NOT exceed the equivalent of 1.0 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products at planting and 0.4 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.
5)	For leafy vegetables make no more than one soil application per crop year.
6)	For spinach a total of 2 shank applications may be made per crop year.
7)	Pre-harvest Interval (PHI): a) Leafy vegetables (except spinach): 7 days b) Spinach: 3 days only if soil application does not exceed 1.0 lb ai/A/year and foliar application of Ridomil Gold Copper (EPA registration, #100-804) formulation does not exceed 0.25 lb ai mefenoxam or metalaxyl/A/year. Otherwise, the PHI is 21 days for spinach.

7.5 Root and Tuber Vegetables

7.5.1 CROP GROUP 1, EXCEPT CARROT, GINSENG, POTATO AND SUGAR BEET

Crops (including all cultivars, varieties, and/or hybrids of these)		
Arracacha	Chervil, turnip-rooted	Rutabaga
Arrowroot	Chicory	Salsify (oyster plant)
Artichoke, Chinese	Chufa	Salsify, black
Artichoke, Jerusalem	Dasheen (taro)	Salsify, Spanish
Burdock, edible	Ginger	Skirret
Canna, edible	Horseradish	Sweet potato
Cassava, bitter	Leren	Tanier (cocoyam)
Cassava, sweet	Parsley, turnip-rooted	Turmeric
Celeriac (celery root)	Parsnip	Turnip
Chayote (root)	Radish	Yam bean (jicama, maniois pea)
	Radish, oriental (daikon)	Yam, true

continued...

7.5.1 CROP GROUP 1, EXCEPT CARROT, GINSENG, POTATO AND SUGAR BEET (continued)

Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Pythium root rot (<i>Pythium</i> spp.) Phytophthora root rot (<i>Phytophthora</i> spp.)	1.0 – 2.0 (0.5-1.0)	Preplant incorporated At planting	Preplant incorporated (broadcast or band): Apply in water or liquid fertilizer and mechanically incorporate in the top 2 inches of soil. Use sufficient water to provide uniform coverage of soil. Soil spray (broadcast or band): Apply in water or liquid fertilizer at planting. For banded applications, use a 7-inch band.
Resistance Management: • Refer to Section 3.2 .			
USE RESTRICTIONS			
1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: 2.0 pt/A (equivalent to 1.0 lb ai/A mefenoxam) 3) Minimum Application Interval: NA 4) Maximum Annual Application Rate: 2.0 pt/A/year (equivalent to 1.0 lb ai/A/year mefenoxam) a) DO NOT exceed 1.0 lb ai/A/year of mefenoxam- and metalaxyl-containing products. 5) Pre-harvest Interval (PHI): NA			

7.5.2 GINSENG

Crops (including all cultivars, varieties, and/or hybrids)			
Ginseng			
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Phytophthora root rot (<i>Phytophthora cactorum</i>)	0.75 (0.375)	Apply in the spring before plants start growing.	For stand establishment, apply uniformly as a soil drench to the soil surface. Apply in 100-400 gallons of water per acre. Follow with additional applications of Ridomil Gold GR.
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2. 			
USE RESTRICTIONS			
1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: 0.75 pt/A (equivalent to 0.375 lb ai/A mefenoxam) 3) Minimum Application Interval: NA 4) Maximum Annual Application Rate: 0.75 pt/A/year (equivalent to 0.375 lb ai/A/year mefenoxam) a) DO NOT exceed 0.375 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products, and 1.125 lb ai/A/year foliar- applied mefenoxam- and metalaxyl-containing products. 5) DO NOT make more than one application of Orondis Gold B per year. 6) Pre-harvest Interval (PHI): NA			

7.5.3 POTATO

Crops (including all cultivars, varieties, and/or hybrids)			
Potato			
Target Disease	Rate (lb ai)	Application Timing	Use Directions
Pink rot <i>(Phytophthora erythroseptica)</i> Pythium leak Pythium seedling disease <i>(Pythium spp.)</i>	0.42 fl oz/ 1,000 row ft Equivalent to 6.1 fl oz/A on 36 inch row spacing (0.19)	At planting	In-furrow spray: Apply directly over the seed pieces in-furrow as a 6- to 8-inch band prior to row closure or use markout application method (incorporated). Orondis Gold B may be impregnated on dry fertilizer or applied in combination with liquid fertilizers.
For soil directed and other applications, refer to Section 8.5.1			
Resistance Management: • Refer to Section 3.2.			

USE RESTRICTIONS
<p>1) Refer to Section 6.1 for additional product use restrictions.</p> <p>2) Maximum Single Application Rate: 0.42 fl oz/1000 row ft (equivalent to 0.19 lb ai/A mefenoxam)</p> <p>3) Minimum Application Interval: NA</p> <p>4) Maximum Annual Application Rate: 0.42 fl oz/1000 row ft (equivalent to 0.19 lb ai/A/year mefenoxam)</p> <p>a) DO NOT exceed 0.34 lb ai/A/year of soil-applied and 0.40 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.</p> <p>5) DO NOT apply Orondis Gold B beyond the at-planting stage.</p> <p>6) DO NOT use the “dribble” application method.</p> <p>7) Pre-harvest Interval (PHI): NA</p>

7.6 Tobacco

Crops (including all cultivars, varieties, and/or hybrids)			
Tobacco			
Target Disease	Rate	Application Timing	Use Directions
Black shank (<i>Phytophthora parasitica</i> var. <i>nicotianae</i>)	Transplant water: 4 - 8 fl oz/200 gallons water	At transplant of tobacco seedlings. Make at least one subsequent application of Orondis Gold B at first cultivation and/or layby if necessary.	Transplant water: Apply in transplant furrow while planting tobacco seedlings. Apply 4-8 fl oz/A (0.25-0.5 pt/A) in at least 200 gallons of transplant water per acre. Use the highest specified rate if the disease epidemic is expected to be severe. Pre-mixing Orondis Gold B in a tank separate from the transplant water source tank will help to prevent incompatibility with fertilizers or other pesticides in the transplant water solution.

Target Disease	Rate	Application Timing	Use Directions
Black shank (<i>Phytophthora parasitica</i> var. <i>nicotianae</i>) (continued)	Transplant water: 4 - 8 fl oz/200 gallons water	At transplant of tobacco seedlings. Make at least one subsequent application of Orondis Gold B at first cultivation and/or layby if necessary.	Consult local extension bulletins for additional use directions. For best results against black shank, use tobacco varieties that have high resistance to black shank and use crop rotation. In fields with a history of severe black shank, use the highest specified rate and plant variety resistant to the race of <i>Phytophthora</i> present (Burley L8 hybrids are resistant to only <i>Phytophthora</i> Race 0).
For soil directed, refer to Section 8.6			
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2. 			

continued...

7.6 Tobacco (continued)

Precautions:

- There is a risk of plant injury with transplant water application, especially when applied in less than 200 gallons of carrier volume per acre. Crop injury is more likely when applied in less than 100 gallons/acre. Injury is temporary and should disappear within three weeks.
- Failure to adequately control nematodes in fields treated with Orondis Gold B may result in poor control of black shank.
- **DO NOT** apply to stressed seedlings or during hot and dry conditions due to injury potential.

USE RESTRICTIONS

- 1) Refer to **Section 6.1** for additional product use restrictions.
- 2) **Maximum Single Application Rate:** 3.0 pt/A (equivalent to 1.5 lb ai/A mefenoxam)
- 3) **Minimum Application Interval:** 14 days
- 4) **Maximum Annual Application Rate:** 3.0 pt/A/year (equivalent to 1.5 lb ai/A/year mefenoxam)
 - a) **DO NOT** exceed 1.5 lb ai/A/year of mefenoxam- and metalaxyl-containing products.
- 5) **DO NOT** use in high black shank areas on highly susceptible flue-cured varieties.
- 6) **DO NOT** use Orondis Gold B for black shank control in PA.
- 7) **Pre-harvest Interval (PHI):** NA

8.0 CROP USE DIRECTIONS FOR SOIL DIRECTED APPLICATIONS

- The restricted-entry interval (REI) for soil-directed applications is 48 hours.

8.1 Asparagus

Crops (including all cultivars, varieties, and/or hybrids)			
Asparagus			
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Crown rot Spear rot (<i>Phytophthora</i> spp.)	1.0 (0.5)	Cutting beds: Apply 30 to 60 days before the first cutting. Apply again just before the beginning of harvest. New plantings: Apply after planting seedlings or after covering one-year old crowns.	Soil spray (broadcast or band): Apply as a soil spray (broadcast or band).
Resistance Management: <ul style="list-style-type: none">Refer to Section 3.2.			

continued...

8.1 Asparagus (continued)

USE RESTRICTIONS	
1)	Refer to Section 6.1 for additional product use restrictions.
2)	Maximum Single Application Rate: 1.0 pt/A (equivalent to 0.5 lb ai/A mefenoxam)
3)	Minimum Application Interval: 30 days
4)	Maximum Annual Application Rate: 2.0 pt/A/year (equivalent to 1.0 lb ai/A/year mefenoxam)
a)	DO NOT exceed 1.0 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products.
5)	Pre-harvest Interval (PHI): 1 day

8.2 Berry and Small Fruit

8.2.1 BUSHBERRY, CROP SUBGROUP 13-07B

Crops (including all cultivars, varieties, and/or hybrids of these)		
Aronia berry	Elderberry	Jostaberry
Blueberry, highbush	European barberry	Juneberry (Saskatoon Berry)
Blueberry, lowbush	Gooseberry	Lingonberry
Buffalo currant	Highbush cranberry	Native currant
Chilean guava	Honeysuckle, edible	Salal
Currant, black	Huckleberry	Sea buckthorn
Currant, red		

Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Root rot (<i>Phytophthora</i> spp.)	3.6 (1.8)	<p>New Plantings: Apply after initial at-planting application.</p> <p>Established Plantings: Apply before the plants start to grow in the spring.</p>	<p>Soil Application (band or via drip/micro-sprinkler irrigation)</p> <p>New Plantings: Reapply once during a period favorable for root rot.</p> <p>Established Plantings: Make a soil-directed application towards the base of the plant in a 3-ft band over the row or via the drip irrigation. One additional application may be made to coincide with the period most favorable for root rot development.</p> <p>Use Orondis Gold B in conjunction with good cultural practices to minimize disease.</p>
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2. 			
Precaution: <ul style="list-style-type: none"> Orondis Gold B will not revitalize plants showing moderate to severe root rot symptoms. 			

continued...

8.2.1 BUSHBERRY, CROP SUBGROUP 13-07B (continued)

USE RESTRICTIONS	
1)	Refer to Section 6.1 for additional product use restrictions.
2)	Maximum Single Application Rate: 3.6 pt/A (equivalent to 1.8 lb ai/A mefenoxam)
3)	Minimum Application Interval: 90 days
4)	Maximum Annual Application Rate: 7.2 pt/A/year (equivalent to 3.6 lb ai/A/year mefenoxam)
a)	DO NOT exceed 3.6 lb ai/A/year of soil applied mefenoxam- and metalaxyl-containing products.
5)	Pre-harvest Interval (PHI): 0 days

8.2.2 CANEBERRY, CROP SUBGROUP 13-07A

Crops (including all cultivars, varieties, and/or hybrids of these)			
Blackberry Loganberry		Raspberry, Red Raspberry, black	Raspberry, wild
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Root rot (<i>Phytophthora</i> spp.)	3.6 (1.8)	<p>New plantings: After the initial at-planting application (Section 7.1.2), reapply once during a period favorable for root rot.</p> <p>Established plantings: Apply before the plants start to grow in the spring. One additional application may be made to coincide with the period most favorable for root rot development.</p>	<p>Soil application: Apply by band or via drip/microsprinkler irrigation.</p> <p>Established plantings: Make the spring application towards the base of the plant in a 3-ft band over the row or via the drip irrigation. Use Orondis Gold B in conjunction with good cultural practices to minimize disease.</p>
For soil directed or soil incorporated applications, refer to Section 7.1.2			

continued...

8.2.2 CANEBERRY, CROP SUBGROUP 13-07A (continued)

Resistance Management: <ul style="list-style-type: none">• Refer to Section 3.2.
Precaution: <ul style="list-style-type: none">• Orondis Gold B will not revitalize plants showing moderate to severe root rot symptoms.
USE RESTRICTIONS
<ol style="list-style-type: none">1) Refer to Section 6.1 for additional product use restrictions.2) Maximum Single Application Rate: 3.6 pt/A (equivalent to 1.8 lb ai/A mefenoxam)3) Minimum Application Interval: 90 days4) Maximum Annual Application Rate: 7.2 pt/A/year (equivalent to 3.6 lb ai/A/year mefenoxam)<ol style="list-style-type: none">a) DO NOT exceed 3.6 lb ai/A/year of soil-applied and 0.2 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.5) DO NOT use an adjuvant.6) Pre-harvest Interval (PHI): 45 days

8.3 Cucurbit Vegetables, Crop Group 9

Crops (including all cultivars, varieties, and/or hybrids of these)			
Chayote (fruit)	Momordica spp.	Mango melon	Squash, summer
Chinese waxgourd	Balsam apple	Persian melon	Zucchini
(Chinese preserving melon)	Balsam pear	Pineapple melon	Squash, winter
Citron melon	Bittermelon	Santa Claus melon	Acorn squash
Cucumber	Chinese cucumber	Snake melon	Butternut squash
Gherkin	Muskmelon	True cantaloupe	Calabaza
Gourd, edible	Cantaloupe	Pumpkin	Hubbard squash
Hyotan	Casaba	Squash, summer	Spaghetti squash
Cucuzza	Crenshaw melon	Crookneck squash	Watermelon
Hechima	Golden pershaw melon	Scallop squash	
Chinese okra	Honeydew melon	Straightneck squash	
	Honey balls	Vegetable marrow	

continued...

8.3 Cucurbit Vegetables, Crop Group 9 (continued)

Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Root rot (<i>Pythium</i> spp) Suppression: Phytophthora blight (<i>Phytophthora capsici</i>)	0.25 – 0.4 (0.125- 0.20)	If soil applications were made at planting, two additional applications may be made at 20- to 30-day intervals.	For soil spray (directed) applications, direct the spray to the base of the plants and cover 6-8 inches of the soil on either side of the plants. Incorporate mechanically or sprinkler-irrigate to move the Orondis Gold B into the root zone. Injection (drip irrigation): Inject Orondis Gold B into the irrigation water at the labeled rates.
For soil-injected or soil incorporated applications, refer to Section 7.2 .			
Resistance Management: • Refer to Section 3.2 .			

USE RESTRICTIONS
<p>1) Refer to Section 6.1 for additional product use restrictions.</p> <p>2) Maximum Single Application Rate: 0.4 pt/A (equivalent to 0.2 lb ai/A mefenoxam)</p> <p>3) Minimum Application Interval: 20 days</p> <p>4) Maximum Annual Application Rate: Soil: 2.0 pt/A/year (equivalent to 1.0 lb ai/A/year mefenoxam)</p> <p>a) DO NOT exceed 1.0 lb ai/A/year of soil-applied and 0.5 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.</p> <p>5) Pre-harvest Interval (PHI): 5 days</p>

8.4 Fruiting Vegetables

8.4.1 CROP GROUP 8, EXCEPT TOMATO

Crops (including all cultivars, varieties, and/or hybrids of these)			
Eggplant Groundcherry Pepino	Pepper Bell Chili Cooking	Pepper Pimento Sweet Tomatillo	
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Crown rot (<i>Phytophthora capsici</i>)	1.0 (0.5)	Make 2 post-directed applications at 30-day intervals following transplanting. Apply before plants are infected to obtain satisfactory results.	For banded spray application , direct the spray to the base of the plants and cover 6-8 inches of the soil on either side of the plants. Incorporate mechanically or sprinkler-irrigate to move the Orondis Gold B into the root zone. For shank application , apply in liquid fertilizer, shanked in as a banded treatment to either side of the plant. For Injection (drip irrigation) , inject Orondis Gold B into the irrigation water.

For soil-injected or soil incorporated applications, refer to Section 7.3.1 .
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2.
Precautions: <ul style="list-style-type: none"> Application of Orondis Gold B may cause some yellowing of pepper leaves. Plants already infected with <i>Phytophthora capsici</i> cannot be cured with Orondis Gold B. The foliar blight phase of <i>Phytophthora</i> cannot be cured with Orondis Gold B.
USE RESTRICTIONS
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: 1.0 pt/A (equivalent to 0.5 lb ai/A mefenoxam) Minimum Application Interval: 30 days Maximum Annual Application Rate: 2.0 pt/A/year (equivalent to 1.0 lb ai/A/year mefenoxam) <ol style="list-style-type: none"> DO NOT exceed 1.5 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products and 0.5 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products. Pre-harvest Interval (PHI): 7 days

8.4.2 TOMATO

Crops (including all cultivars, varieties, and/or hybrids)			
Tomato			
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Fruit rot (<i>Phytophthora</i> spp.) Root rot (<i>Pythium</i> spp.)	1.0 (0.5)	4-6 weeks after planting. If needed, make a second application up to 1 week before harvest.	Soil spray (broadcast or band) or soil injection: Apply as a directed soil surface spray under the vines or injected into the beds with water or liquid fertilizer.
For soil-injected or soil incorporated applications, refer to Section 7.3.2 .			
Resistance Management: <ul style="list-style-type: none">• Refer to Section 3.2.			

USE RESTRICTIONS
<ol style="list-style-type: none">1) Refer to Section 6.1 for additional product use restrictions.2) Maximum Single Application Rate: 1.0 pt/A (equivalent to 0.5 lb ai/A mefenoxam)3) Minimum Application Interval: 28 days4) Maximum Annual Application Rate: 3.0 pt/A/year (equivalent to 1.5 lb ai/A/year mefenoxam)<ol style="list-style-type: none">a) DO NOT exceed 1.5 lb ai/A/year of soil-applied and 0.5 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.5) Pre-harvest Interval (PHI): 7 days

8.5 Root and Tuber Vegetables

8.5.1 POTATO

Crops (including all cultivars, varieties, and/or hybrids)			
Potato			
Target Disease	Rate fl oz/A (lb ai)	Application Timing	Use Directions
Storage rots Pink rot (<i>Phytophthora erythrosepica</i>) Pythium leak (<i>Pythium</i> spp.)	3.2 (0.1)	At tuber initiation when the largest tubers are the size in diameter of a nickel. Usually coincides with initiation of flowering. Make a second application 14 days later and, if the field has a history of storage rot problems, a third application 14 days after the second application.	Make broadcast or soil-directed applications by air, ground, or chemigation. If the field has a history of storage rot problems, make a third application. If foliar diseases are expected, then a tank mix with a labeled rate of mancozeb or chlorothalonil products is required. Use in conjunction with other management practices such as crop rotation and resistant varieties.
For soil-injected or soil incorporated applications, refer to Section 7.5.3 .			
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2. 			

USE RESTRICTIONS
<ol style="list-style-type: none">1) Refer to Section 6.1 for additional product use restrictions.2) Maximum Single Application Rate: 3.2 fl oz/A (equivalent to 0.1 lb ai/A mefenoxam)3) Minimum Application Interval: 14 days4) Maximum Annual Application Rate: 9.6 fl oz/A/year (equivalent to 0.3 lb ai/A/year mefenoxam)<ol style="list-style-type: none">a) DO NOT exceed 0.34 lb ai/A/year of soil-applied and 0.40 lb ai/A/year of foliar-applied mefenoxam- and metalaxyl-containing products.5) Pre-harvest Interval (PHI): 14 days

8.6 Tobacco

Crops (including all cultivars, varieties, and/or hybrids)			
Tobacco			
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Black shank (<i>Phytophthora parasitica</i> var. <i>nicotianae</i>)	1.0 – 2.0 (0.5-1.0) For no-till tobacco: 0.50 – 1.0 (0.25-0.5)	One application at lay-by or one application at the first cultivation and a second application at lay-by. Apply preventatively for effective black shank control. If black shank is expected early in the season, apply as near as possible to transplanting followed by sequential applications.	Soil spray (broadcast or band): Position the nozzles so the spray is deposited under the plants and covered with soil by the cultivator. Use the higher specified rate if the disease epidemic is expected to be severe. Consult local extension bulletins for additional use directions. For best results against black shank, use tobacco varieties that have high resistance to black shank and use crop rotation. In fields with a history of severe black shank, use the highest specified rate and plant variety resistant to the race of <i>Phytophthora</i> present (Burley L8 hybrids are resistant to only <i>Phytophthora</i> Race 0).

For soil-injected or soil incorporated applications, refer to Section 7.6 .
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2.
Precaution: <ul style="list-style-type: none"> Failure to adequately control nematodes in fields treated with Orondis Gold B may result in poor control of black shank.
USE RESTRICTIONS
1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: 2.0 pt/A (equivalent to 1.0 lb ai/A mefenoxam) 3) Minimum Application Interval: 14 days 4) Maximum Annual Application Rate: 3.0 pt/A/year (equivalent to 1.5 lb ai/A/year mefenoxam) a) DO NOT exceed 1.5 lb ai/A/year of mefenoxam- and metalaxyl-containing products. 5) DO NOT use in high black shank areas on highly susceptible flue-cured varieties. 6) DO NOT use Orondis Gold B for black shank control in PA. 7) Pre-harvest Interval (PHI): NA

9.0 STORAGE AND DISPOSAL

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Handling (less than or equal to 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

continued...

Container Handling (greater than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

10.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT**

LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Orondis®, Ridomil Gold®, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



Viton® is a registered trademark of E.I. DuPont de Nemours & Company, Inc.

©2021 Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1202B-L1H 1120
4134440

MEFENOXAM GROUP 4 FUNGICIDE

Orondis® Gold B Fungicide

For the control of certain diseases in listed crops caused by the Oomycete class of fungi

Active Ingredient:	
Mefenoxam*	45.3%
Other Ingredients:	54.7%
Total:	100.0%

*CAS No. 70630-17-0 and 69516-34-3

Orondis® Gold B is formulated as a soluble concentrate and contains 4 lb active ingredient per gallon.

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1202 EPA Est. 072344-MO-004

2.5 quarts
(80 fluid ounces) Net Contents

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

KEEP OUT OF REACH OF CHILDREN. CAUTION

FIRST AID If in **eyes**: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on **skin or clothing**: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If **swallowed**: Call a poison control center or doctor immediately for treatment

advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by the poison control center or doctor. Do not give anything to an unconscious person. If **inhaled**: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOTLINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call **1-800-888-8372**.

Orondis® and the SYNGENTA Logo are trademarks of a Syngenta Group Company.

SCP 1202B-L1H 1120 4134440

syngenta®

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: Mefenoxam is known to leach through the soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a

high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

Physical or Chemical Hazards: Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according

to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Manufactured for: Syngenta Crop Protection, LLC

P.O. Box 18300

Greensboro, NC 27419-8300

SCP 1202B-L2E 1120

4148494

