

AZOXYSTROBIN GROUP 11 FUNGICIDE

FLUDIOXONIL GROUP 12 FUNGICIDE

PULL HERE TO OPEN ►

DIFENOCONAZOLE GROUP 3 FUNGICIDE



Stadium[®]

Fungicide

syngenta[®]

Active Ingredients:

Azoxystrobin*	12.51%
Fludioxonil**	12.51%
Difenoconazole***	9.76%

Other Ingredients:	65.22%
--------------------	--------

Total:	100.0%
--------	--------

*CAS No. 131860-33-8

**CAS No. 131341-86-1

***CAS No. 119446-68-3

Stadium Fungicide is formulated as a suspension concentrate and contains 1.197 pounds of azoxystrobin, 1.197 pounds of fludioxonil, and 0.934 pounds of difenoconazole per gallon.

KEEP OUT OF REACH OF CHILDREN. CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1453 EPA Est. 100-NE-001

SCP 1453A-L1B 0823

4191516

Product ID. **51616**

5 gallons
Net Contents

[®]

TABLE OF CONTENTS

1.0 FIRST AID

PRECAUTIONARY STATEMENTS

2.0 PRECAUTIONARY STATEMENTS

- 2.1 Hazards to Humans and Domestic Animals
- 2.2 Personal Protective Equipment (PPE)
- 2.3 User Safety Requirements
- 2.4 User Safety Recommendations
- 2.5 Environmental Hazards
 - 2.5.1 GROUNDWATER ADVISORY
- 2.6 Physical or Chemical Hazards

DIRECTIONS FOR USE

3.0 PRODUCT INFORMATION

- 3.1 Resistance Management
- 3.2 Integrated Pest Management (IPM)

4.0 APPLICATION DIRECTIONS

- 4.1 Methods of Application
- 4.2 Application Equipment
- 4.3 Application Volume and Spray Coverage
- 4.4 Mixing Directions
 - 4.4.1 STADIUM FUNGICIDE ALONE
 - 4.4.2 TANK-MIX PRECAUTIONS
 - 4.4.3 TANK-MIX COMPATIBILITY
 - 4.4.4 STADIUM FUNGICIDE IN TANK MIXTURES

5.0 POST-HARVEST CROP USE DIRECTIONS

- 5.1 Tuberous and Corm Vegetables Subgroup 1C

6.0 STORAGE AND DISPOSAL

7.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

1.0 FIRST AID

FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOTLINE NUMBER For 24-hour medical emergency assistance (human or animal) or chemical emergency assistance (spill, leak, fire, or accident), call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

CAUTION/PRECAUCIÓN

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

2.2 Personal Protective Equipment (PPE)

Mixers, loaders, applicators and all other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neo-prene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, and Viton™ ≥ 14 mils

2.3 User Safety Requirements

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.4 User Safety Recommendations

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

2.5 Environmental Hazards

This pesticide is toxic to fish, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic estuarine/marine organisms in water adjacent to treated area.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

2.5.1 GROUNDWATER ADVISORY

These chemicals have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

2.6 Physical or Chemical Hazards

Do not use or store near heat or open flame. Do not store near or use with oxidizing agents.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not formulate this product into other end-use products.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

3.0 PRODUCT INFORMATION

Stadium Fungicide is a broad-spectrum fungicide with systemic and curative properties for the control of many important plant diseases in post-harvest treatment facilities. All applications must be made according to the use directions that follow.

3.1 Resistance Management

AZOXYSTROBIN	GROUP	11	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE

Stadium Fungicide contains three fungicides with different modes of action. Azoxystrobin is a strobilurin in Group 11. Fludioxonil is a phenylpyrrole in Group 12. Difenoconazole is a triazole fungicide in Group 3. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Syngenta Crop Protection encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label. Stadium Fungicide should not be alternated or tank-mixed with any fungicide to which resistance has already developed. If resistant isolates to Group 3, 11 or 12 fungicides are present, efficacy can be reduced.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of azoxystrobin or other Group 11 fungicides, fludioxonil or other Group 12 fungicides, and difenoconazole or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crop and pathogens.
- For further information or to report suspected resistance contact Syngenta at 1-866-Syngent(a) (866-796-4368). You can also contact your pesticide distributor or university extension specialist to report resistance.

3.2 Integrated Pest Management (IPM)

Stadium Fungicide should be integrated into an overall disease and pest management strategy (IPM) whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area.

4.0 APPLICATION DIRECTIONS

4.1 Methods of Application

Use Stadium Fungicide as an in-line spray for the control of certain post-harvest rots caused by Silver Scurf (*Helminthosporium solani*) and *Fusarium* species.

4.2 Application Equipment

- All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.
- Use T-jet, CDA or similar application system for in-line aqueous application.

4.3 Application Volume and Spray Coverage

- Ensure proper coverage from the in-line aqueous spray.
- Ensure the tubers are not excessively wet as efficacy will be impacted.

4.4 Mixing Directions

1. Thoroughly clean spray equipment before using this product.
2. Prepare no more spray mixture than is needed for the immediate operation.
3. Keep product container tightly closed when not in use.
4. Agitate the spray solution before and during application.
5. Do not let the spray mixture stand overnight in the spray tank.
6. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

4.4.1 STADIUM FUNGICIDE ALONE

1. Prepare no more spray mixture than is required for the immediate operation.
2. Thoroughly clean spray equipment before using this product.
3. Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
4. With the agitator running, add Stadium Fungicide to the tank.
5. Continue agitation while adding the remainder of the water.
6. Begin application of the spray solution after Stadium Fungicide has completely dispersed into the mix water.
7. Maintain agitation until all of the mixture has been sprayed.
8. Rinse spray tank thoroughly with clean water after each day's use.

4.4.2 TANK-MIX PRECAUTIONS

- Plant tolerance has been found acceptable for all crops on the label; however, not all possible tank-mix combinations have been tested under all conditions. When possible, it is recommended to test the combinations on a small portion of the crop to ensure a phytotoxic response will not occur as a result of application.
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitation and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Tank mixes of Stadium Fungicide with other pesticides, or any other additives not specifically labelled for use with Stadium Fungicide may result in tank mix incompatibility or unsatisfactory performance. In such cases, always check tank mix compatibility by conducting a jar test according to guidance in **Section 4.4.3** before actual tank mixing.

4.4.3 TANK-MIX COMPATIBILITY

- Conduct a jar test using a 1 pt to 1 qt container with lid by adding water or other intended carrier to the jar.
- Next, add the appropriate amount of pesticide(s) or tank mix partner(s) in their relative proportions based on recommended label rates. Add tank mix components separately in the order described in the tank-mixing section, **Section 4.4.4**. After each addition, shake or stir gently to thoroughly mix.
- After all ingredients have been added, put the lid on the jar, tighten and invert the jar 10 times to mix.
- After mixing, let the mixture stand 15 – 30 minutes and then examine for signs of incompatibility such as obvious separation, large flakes, precipitates, gels or heavy oily film on the jar.
- If the mixture remains mixed or can be remixed readily, it is physically compatible and can be used.
- If the mixture is incompatible, repeat the test using a compatibility agent at the recommended rate. Or, if applicable, slurry dry formulations in water before adding to the jar. If incompatibility is still observed after following these procedures, do not use the mixture.
- After compatibility testing is complete, dispose of any pesticide wastes in accordance with the storage and disposal section, (**Section 6.0**) of this label.

4.4.4 STADIUM FUNGICIDE IN TANK MIXTURES

1. Prepare no more spray mixture than is required for the immediate operation.
2. Thoroughly clean spray equipment before using this product.
3. Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
4. With the agitator running, add the tank-mix partner(s) into the tank.
5. Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Stadium Fungicide to the spray tank.
6. Allow Stadium Fungicide to completely disperse.
7. Spray the mixture with the agitator running.
8. Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix product label.
9. Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed.
10. This product must not be mixed with any product which prohibits such mixing.
11. Rinse spray tank thoroughly with clean water after each day's use.

5.0 POST-HARVEST CROP USE DIRECTIONS

5.1 Tuberous and Corm Vegetables Subgroup 1C

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Arracacha	Cassava, Sweet	Potato	
Arrowroot	Chayote (root)	Potato, Sweet	
Artichoke, Chinese	Chufa,	Tanier	
Artichoke, Jerusalem	Dasheen	Turmeric	
Canna, Edible	Ginger	Yam Bean	
Cassava, Bitter	Leren	Yam, True	
Target Disease	Rate (fl oz)	Application Timing	Use Directions
Silver Scurf Fusarium Dry Rot	1.0 fl oz/ 2000 lb of tubers	In-Line Aqueous Spray	Ensure proper coverage of the tubers. Tubers should be tumbling as they are treated. Mix the fungicide solution in an appropriate amount of water for the crop being treated. Use T-jet, CDA, or similar application system.
Resistance Management: • Refer to Section 3.1.			
USE RESTRICTIONS			
1) Maximum Rate per Application: 1.0 fl oz/2000 lb of tubers 2) Maximum Rate per 2000 lb of Tubers: DO NOT exceed 0.0094 lb ai of azoxystrobin. DO NOT exceed 0.0094 lb ai of fludioxonil. DO NOT exceed 0.0073 lb ai of difenoconazole. 3) DO NOT make more than one post-harvest application to the tubers.			

6.0 STORAGE AND DISPOSAL

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in the original container in cool, dry, secure place.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling (less than or equal to 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to contain spills, leaks, and other accidents to prevent further exposure of facilities and equipment. Absorb spilled product with absorbing materials and dispose of in an approved waste disposal facility. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

7.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Stadium®, the ALLIANCE FRAME
the Syngenta Logo and the PURPOSE ICON
are trademarks of a Syngenta Group Company

Viton™ is a trademark of The Chemours Company FC, LLC

©2023 Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 1453A-L1B 0823
4191516**

AZOXYSTROBIN	GROUP	11	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE

Stadium® Fungicide

Active Ingredients:

Azoxystrobin*	12.51%
Fludioxonil**	12.51%
Difenoconazole***	9.76%
Other Ingredients:	65.22%
Total:	100.0%

*CAS No. 131860-33-8

**CAS No. 131341-86-1

***CAS No. 119446-68-3

Stadium Fungicide is formulated as a suspension concentrate and contains 1.197 pounds of azoxystrobin, 1.197 pounds of fludioxonil, and 0.934 pounds of difenoconazole per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION/ PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1453

EPA Est. 100-NE-001

Product ID. **51616**

5 gallons
Net Contents

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOTLINE NUMBER: For 24-hour medical emergency assistance (human or animal) or chemical emergency assistance (spill, leak, fire, or accident), call **1-800-888-8372**.

Precautionary Statements

Hazards to Humans and Domestic Animals CAUTION/PRECAUCIÓN

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Environmental Hazards: This pesticide is toxic to fish, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic estuarine/marine organisms in water adjacent to treated area.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: These chemicals have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the

permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

Physical or Chemical Hazards: Do not use or store near heat or open flame. Do not store near or use with oxidizing agents.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in the original container in cool, dry, secure place.

Pesticide Disposal: Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling (less than or equal to 5 gallons): Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Stadium® and the Syngenta logo are trademarks of a Syngenta Group Company

©2023 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 1453A-L1B 0823
4191516**

syngenta®

