




Ultimate control of downy mildews

Orondis[®] Ultra combines two products, Orondis (oxathiapiprolin) and Revus[®] (mandipropamid) fungicides, to aid growers in hitting an out-of-the-park home run against downy mildews in leafy and brassica vegetables. As the only active ingredient in FRAC Group 49, oxathiapiprolin is not cross-resistant to any other fungicide.

- Outstanding preventive and residual disease control
- Systemic, translaminar movement and redistribution to protect developing leaves
- Rainfast within 30 minutes after spray deposits have dried
- Excellent crop safety
- Foundation fungicide product in spray programs with other modes of action
- Built-in resistance management

ROBUST CONTROL OF LETTUCE DOWNY MILDEW IN FUNGICIDE PROGRAM

Orondis Ultra in a fungicide program shows superior downy mildew control in lettuce



Source: Trial USWB0F1182016/Crop Science Services/CA/2016




Commercial reference program




Orondis Ultra program

SUPERIOR SPINACH DOWNY MILDEW CONTROL



OUTSTANDING CONTROL OF BROCCOLI FOLIAR DOWNY MILDEW



SYSTEMIC DOWNY MILDEW CONTROL IN BROCCOLI



ORONDIS ULTRA LABEL AT A GLANCE*

Source: Trial USVJ0F1222016/FL/2016

	Leafy greens (Crop subgroup 4A)	Brassica vegetables (Crop subgroup 5A)
Spectrum of activity	Downy mildew (<i>Bremia lactucae</i>) - lettuce, Downy mildew (<i>Peronospora farinosa</i>) - spinach	Downy mildew (<i>Peronospora parasitica</i>)
Rate	5.5 - 8.0 fl oz/A	
Maximum amount per season	32 fl oz/A	
Minimum gallons per acre (GPA)	Ground: 15, Aerial: 5	
Adjuvants	A spreading/penetrating adjuvant** at labeled agricultural use rates may enhance disease control	A spreading/penetrating adjuvant** must be added at labeled agricultural use rates
Reentry interval (REI)	4 hours	
Preharvest interval (PHI)	1 day	
Crop rotation restrictions	0 days to crops on label; 30 days to cereals and grass animal feeds; 180 days to all other crops	

* Always consult the product label for complete use directions and application information.

** Spreading/penetrating types of adjuvant include non-ionic surfactants, organosilicones, or blends.

Photos are either the property of Syngenta or used under agreement.

©2017 Syngenta: **Always read and follow label instructions. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.** Actigard®, Orondis®, Revus®, the Alliance Frame, the Purpose Icon and the Syngenta logo are trademarks of a Syngenta Group Company. Zampro® is a registered trademark of BASF. Fosphite® is a registered trademark of JH BioTech, Inc. Manzate® is a registered trademark of DuPont or its affiliates. ProPhyt® is a trademark of Helena Chemical Company. Ranman® is a trademark of FMC Corp. Other trademarks displayed or otherwise used herein are the property of a Syngenta Group Company or the respective trademark owners.