



## FOR RELIABLE, EASY BROAD-SPECTRUM INSECT CONTROL

For a modern insecticide that provides reliable, broad-spectrum control of damaging pests, trust Voliam Flexi<sup>®</sup> insecticide. Voliam Flexi gives growers the long-lasting residual they need to control key sucking, chewing and lepidopteran pests in citrus and tree fruit. With dual modes of action and a convenient premix formulation, Voliam Flexi is the easy way to protect your crops and your bottom line.

### THE VOLIAM FLEXI ADVANTAGE

- Complementary, dual modes of action result in convenient broad-spectrum control of key sucking and chewing pests
- An innovative mode of action for lepidopteran control to assist with resistance management
- Excellent residual control and longer interval between applications
- Removes the hassle of tank mixing: saves time, reduces exposure, reduces chances for mistakes, fewer calculations
- Easier compliance – one label has all the information needed to meet federal and local regulations
- Clear, concise MRL information readily available for export markets

### VOLIAM FLEXI USE RECOMMENDATIONS\*

#### Citrus, Pome Fruit and Stone Fruit

- Apply before pests reach damaging levels
- When listed, use higher rate for heavy infestations
- Use sufficient water for thorough coverage
- For best results, apply in 100-150 gallons of water per acre
- Do not apply dilute applications of more than 200 GPA
- Use of an adjuvant is permitted on citrus, pome and stone fruit
- REI = 12 hours

\*Always consult the Voliam Flexi label for a complete list of registered crops, use directions and application information

#### Citrus

- Allow seven days between applications
- Do not exceed 14 oz/A/season of Voliam Flexi
- Application is permitted by ground and air; aerial application may result in slower activity and reduced control
- PHI = 1 day

#### Pome Fruit

- Allow 10 days between applications
- Do not exceed 16 oz/A/season or four applications per season of Voliam Flexi
- Do not apply by air
- PHI = 35 days

#### Stone Fruit

- Allow 10 days between applications
- Do not exceed 14 oz/A/season or three applications per season of Voliam Flexi
- Do not apply by air
- PHI = 14 days



## VOLIAM FLEXI LABEL AT A GLANCE\*

| Crop         | Pests  | Rate Per Application | Preharvest Interval (PHI) | Maximum Per Growing Season  | Water Volume   |
|--------------|--|----------------------|---------------------------|---|--|
| Citrus Fruit | Aphids, Leafhoppers  | 4-5 oz/acre          | 1 Day                     | 14 oz/acre (or 0.172 lb a.i. of thiamethoxam-containing products or 0.2 lb a.i. of chlorantraniliprole-containing products per acre per growing season) | Do not use less than 30 GPA for ground applications or 5 GPA for aerial applications |
|              | Asian citrus psyllid, Citrus black fly, Citrus leafminer, Mealybugs, Sharpshooters, Soft scales, Whiteflies  | 5-7 oz/acre          |                           |   |  |
|              | Ants (except carpenter, fire, harvester and pharaoh ants), Armored scales, Citrus peelminer, Citrus root weevil adults, Crickets, Fruit fly, Grasshoppers, Katydid, Plant bugs, Stink bugs, Thrips (foliage feeding) | 6-7 oz/acre          |                           |   |  |
| Pome Fruit   | Codling moth, Green fruitworm, Leafhoppers, Obliquebanded leafroller, Oriental fruit moth, Spotted tentiform leafminer, Tufted apple budmoth   | 4-7 oz/acre          | 35 Days                   | 16 oz/acre (or 0.25 lb a.i. of thiamethoxam-containing products or 0.2 lb a.i. of chlorantraniliprole-containing products per acre per growing season)  | Do not use less than 50 GPA for ground applications                                  |
|              | Apple aphid, Apple grain aphid, Comstock mealybug, European apple sawfly, Green peach aphid, Leafminers, Mullein bug, Plum curculio, Rosy apple aphid  | 6-7 oz/acre          |                           |   |  |
|              | Pear psylla  | 7 oz/acre            |                           |   |  |
| Stone Fruit  | Aphids, Codling moth, Green fruitworm, Leafhoppers, Oriental fruit moth, Peach twig borer  | 4-7 oz/acre          | 14 Days                   | 14 oz/acre (or 0.172 lb a.i. of thiamethoxam-containing products or 0.2 lb a.i. of chlorantraniliprole-containing products per acre per growing season) | Do not use less than 30 GPA for ground applications                                  |
|              | Cherry fruit fly, Plum curculio, Stink bugs, Tarnished plant bug, Thrips   | 6-7 oz/acre          |                           |   |  |

\*Always consult the Voliam Flexi label for a complete list of registered crops, use directions and application information.

For more information, visit [www.syngentacropprotection.com](http://www.syngentacropprotection.com), [www.FarmAssist.com](http://www.FarmAssist.com) or call the Syngenta Customer Center at 1-866-SYNGENT(A) (796-4368).

©2011 Syngenta Crop Protection, LLC, 410 Swing Road, Greensboro, NC 27409.

**Important: Always read and follow label instructions before buying or using Syngenta products. The instructions contain important conditions of sale, including limitations of warranty and remedy. Voliam Flexi is not currently registered for sale or use in all states. Please check with your state or local extension service before buying or using this product.**

Voliam Flexi is highly toxic to bees exposed to direct treatment on blooming crops and weeds. Do not apply this product or allow it to drift onto blooming plants while bees are foraging adjacent to the treatment area. Voliam Flexi<sup>®</sup>, the Purpose icon and the Syngenta logo are trademarks of a Syngenta Group Company.